

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Related Applications

Reference

Background of the invention

Blood vessels are the means by which oxygen and nutrients are supplied to living tissues and waste products are removed from living tissue. Angiogenesis refers to the process by which new blood vessels are formed. See, for example, the review by Folkman and Shing, *J. Biol. Chem.* 267(16):10931-10934, 1992. Thus, where appropriate, angiogenesis is a critical biological process. It is essential in reproduction, development and wound repair. However, inappropriate angiogenesis can have severe negative consequences. For example, it is only after many solid tumors are vascularized as a result of angiogenesis that the tumors have a sufficient supply of oxygen and nutrients that permit it to grow rapidly and metastasize. Because maintaining the rate of angiogenesis in its proper equilibrium is so critical to a range of functions, it must be carefully regulated in order to maintain health. The angiogenesis process is believed to begin with the degradation of the basement membrane by proteases secreted from endothelial cells (EC) activated by mitogens such as vascular endothelial growth factor (VEGF) and basic fibroblast growth factor (bFGF). The cells migrate and proliferate, leading to the formation of solid endothelial cell sprouts into the stromal space, then, vascular loops are formed and capillary tubes develop with formation of tight junctions and deposition of new basement membrane.

Endothelial cells is typically low compared to other

25 tight junctions and deposition of new basement membrane.
In adults, the proliferation rate of endothelial cells is typically low compared to other
cell types in the body. The turnover time of these cells can exceed one thousand days.
Physiological exceptions in which angiogenesis results in rapid proliferation typically occurs
under tight regulation, such as found in the female reproduction system and during wound
30 healing.

The rate of angiogenesis involves a change in the local equilibrium between positive and negative regulators of the growth of microvessels. The therapeutic implications of angiogenic growth factors were first described by Folkman and colleagues over two decades ago (Folkman, *N. Engl. J. Med.* 285:1182-1186, 1971). Abnormal angiogenesis occurs when the body loses at least some control of angiogenesis, resulting in either excessive or insufficient blood vessel growth. For instance, conditions such as ulcers, strokes, and heart attacks may result from the absence of angiogenesis normally required for natural healing. In contrast, excessive blood vessel proliferation can result in tumor growth, tumor spread, blindness, psoriasis and rheumatoid arthritis.

There are instances where a greater degree of angiogenesis is desirable, e.g., increasing blood circulation, wound healing, and ulcer healing. For example, recent investigations have established the feasibility of using recombinant angiogenic growth factors, such as fibroblast growth factor (FGF) family (Yanagisawa-Miwa et al., *Science*, 257:1401-1403, 1992; Baffour et al., *J. Vasc. Surg.* 16:181-91, 1992), endothelial cell growth factor (ECGF; Pu et al., *J. Surg. Res.* 54:575-83, 1993), and more recently, vascular endothelial growth factor (VEGF) to expedite and/or augment collateral artery development in animal models of myocardial and hindlimb ischemia (Takeshita et al., *Circulation*, 90:228-234, 1994; Takeshita et al., *J. Clin. Invest.* 93:662-70, 1994).

Conversely, there are instances where inhibition of angiogenesis is desirable. For example, many diseases are driven by persistent unregulated angiogenesis, also sometimes referred to as "neovascularization". In arthritis, new capillary blood vessels invade the joint and destroy cartilage. In diabetes, new capillaries invade the vitreous of the eye, bleed, and cause blindness. Ocular neovascularization is the most common cause of blindness. Tumor growth and metastasis are angiogenesis-dependent. A tumor must continuously stimulate the growth of new capillary blood vessels for the tumor itself to grow.

The current approved treatment of these diseases is inadequate. Agents which prevent continued angiogenesis, such as drugs (e.g. TNP-470), monoclonal antibodies, antisense nucleic acids and proteins (e.g., angiostatin and endostatin) are currently being tested, but have not been approved. Although preliminary results with the antiangiogenic proteins are promising, they are relatively large in size and are difficult to use and produce. Moreover, proteins are subject to enzymatic degradation. Thus, new agents that inhibit angiogenesis are needed. New antiangiogenic agents that show improvement in size, ease of production, stability and/or potency would be desirable.

Summary of the Invention

It has now been discovered that nucleic acid molecules, including oligonucleotides, have intrinsic antiangiogenesis properties apart from the proteins such nucleic acids may encode.

5 According to one aspect of the invention, methods for inhibiting angiogenesis are provided. The methods include administering to a subject in need of such treatment at least one antiangiogenic nucleic acid molecule in an amount effective to inhibit angiogenesis in the subject. In some embodiments, two or more antiangiogenic nucleic acid molecules are administered. In other embodiments, non-nucleic acid antiangiogenic agents also are administered and agents that are effective against other aspects of an angiogenic condition (e.g., anticancer agents) can also be administered. In some embodiments, the angiogenesis is associated with a condition selected from the group consisting of rheumatoid arthritis, psoriasis, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, Osler-Webber Syndrome, myocardial angiogenesis, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma, and wound granulation. In other embodiments, the angiogenesis is not associated with a cancer or tumor, but may be associated with an eye or ocular disorder such as those described herein. In still other embodiments, the angiogenesis is associated with embryo implantation. In certain embodiments, the angiogenesis is associated with conditions involving excessive or abnormal stimulation of endothelial cells such as but not limited to intestinal adhesions, atherosclerosis, scleroderma, and hypertrophic scars, i.e., keloids.

20 In other aspects of the invention, compositions are provided that include at least one antiangiogenic nucleic acid molecule, formulated in a pharmaceutically-acceptable carrier and in an effective amount for inhibiting angiogenesis. The compositions in certain embodiments include non-nucleic acid antiangiogenic agents and/or agents that are effective against other aspects of an angiogenic condition (e.g., anticancer agents).

25 According to still other aspects the invention, kits are provided that include a first container housing at least one antiangiogenic nucleic acid molecule and instructions for administering the antiangiogenic nucleic acid molecule to a subject having unwanted angiogenesis. In certain embodiments, a second container housing at least one non-nucleic acid antiangiogenic agent is also provided. In other embodiments of the foregoing kits, another container housing at least one anticancer agent is provided. In certain embodiments, the instructions relate to administering the antiangiogenic nucleic acid to a subject having a

10017595-15
TOP SECRET

condition that is not cancer or a tumor, and examples of such conditions are listed throughout the specification.

A nucleic acid molecule is an element of each aspect of the invention. Preferred nucleic acid molecules include at least one sequence set forth as SEQ ID NOs: 1-1093. The nucleic acids useful according to the invention are synthetic or natural (isolated) nucleic acids. The nucleic acid may be administered alone or in conjunction with a pharmaceutically-acceptable carrier and optionally other therapeutic agents. In some embodiments the nucleic acid is a CpG nucleic acid, including those having an unmethylated CpG motif, a T-rich nucleic acid, or a poly G nucleic acid.

The nucleic acid in some embodiments has a nucleotide backbone which includes at least one backbone modification, such as a phosphorothioate modification or other phosphate modification. In some embodiments the modified backbone is a peptide modified oligonucleotide backbone. The nucleotide backbone may be chimeric, or the nucleotide backbone is entirely modified.

The nucleic acid can have any length greater than 6 nucleotides, but in some embodiments is between 8 and 100 nucleotide residues in length. In other embodiments the nucleic acid comprises at least 20 nucleotides, at least 24 nucleotides, at least 27, nucleotides, or at least 30 nucleotides. The nucleic acid may be single stranded or double stranded. In some embodiments the nucleic acid is isolated and in other embodiments the nucleic acid may be a synthetic nucleic acid. The antiangiogenic nucleic acids in some instances are not antisense molecules.

The CpG nucleic acid in one embodiment contains at least one unmethylated CpG dinucleotide having a sequence including at least the following formula: 5' X₁ X₂CGX₃ X₄ 3' wherein C is unmethylated, wherein X₁, X₂, X₃, and X₄ are nucleotides. In one embodiment the 5' X₁ X₂CGX₃ X₄ 3' sequence of the CpG nucleic acid is a non-palindromic sequence, and in other embodiments it is a palindromic sequence.

In some embodiments X₁X₂ are nucleotides selected from the group consisting of: GpT, GpG, GpA, ApA, ApT, ApG, CpT, CpA, CpG, TpA, TpT, and TpG; and X₃X₄ are nucleotides selected from the group consisting of: TpT, CpT, ApT, TpG, ApG, CpG, TpC, ApC, CpC, TpA, ApA, and CpA. In other embodiments X₁X₂ are GpA or GpT and X₃X₄ are TpT. In yet other embodiments X₁ or X₂ or both are purines and X₃ or X₄ or both are pyrimidines or X₁X₂ are GpA and X₃ or X₄ or both are pyrimidines. In one embodiment X₂ is a T and X₃ is a pyrimidine.

In other embodiments the CpG nucleic acid has a sequence selected from the group consisting of SEQ ID NO: 1, 3, 4, 14-16, 18-24, 28, 29, 33-46, 49, 50, 52-56, 58, 64-67, 69, 71, 72, 76-87, 90, 91, 93, 94, 96, 98, 102-124, 126-128, 131-133, 136-141, 146-150, 152-153, 155-171, 173-178, 180-186, 188-198, 201, 203-214, 216-220, 223, 224, 227-240, 242-256, 258, 260-265, 270-273, 275, 277-281, 286-287, 292, 295-296, 300, 302, 305-307, 309-312, 314-317, 320-327, 329, 335, 337-341, 343-352, 354, 357, 361-365, 367-369, 373-376, 378-385, 388-392, 394, 395, 399, 401-404, 406-426, 429-433, 434-437, 439, 441-443, 445, 447, 448, 450, 453-456, 460-464, 466-469, 472-475, 477, 478, 480, 483-485, 488, 489, 492, 493, 495-502, 504-505, 507-509, 511, 513-529, 532-541, 543-555, 564-566, 568-576, 578, 580, 599, 601-605, 607-611, 613-615, 617, 619-622, 625-646, 648-650, 653-664, 666-697, 699-706, 708, 709, 711-716, 718-732, 736, 737, 739-744, 746, 747, 749-761, 763, 766-767, 769, 772-779, 781-783, 785-786, 7900792, 798-799, 804-808, 810, 815, 817, 818, 820-832, 835-846, 849-850, 855-859, 862, 865, 872, 874-877, 879-881, 883-885, 888-904, and 909-913.

In some embodiments the T rich nucleic acid is a poly T nucleic acid comprising 5' TTTT 3'. In yet other embodiments the poly T nucleic acid comprises 5' X₁ X₂ TTTT X₃ X₄ 3' wherein X₁, X₂, X₃ and X₄ are nucleotides. In some embodiments X₁X₂ is TT and/or X₃X₄ is TT. In other embodiments X₁X₂ is selected from the group consisting of TA, TG, TC, AT, AA, AG, AC, CT, CC, CA, CG, GT, GG, GA, and GC; and/or X₃X₄ is selected from the group consisting of TA, TG, TC, AT, AA, AG, AC, CT, CC, CA, CG, GT, GG, GA, and GC.

The T rich nucleic acid may have only a single poly T motif or it may have a plurality of poly T nucleic acid motifs. In some embodiments the T rich nucleic acid comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8 T motifs. In other embodiments it comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8 CpG motifs. In some embodiments the plurality of CpG motifs and poly T motifs are interspersed.

In yet other embodiments at least one of the plurality of poly T motifs comprises at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, or at least 9 contiguous T nucleotide residues. In other embodiments the plurality of poly T motifs is at least 3 motifs and wherein at least 3 motifs each comprises at least 3 contiguous T nucleotide residues or the plurality of poly T motifs is at least 4 motifs and wherein the at least 4 motifs each comprises at least 3 contiguous T nucleotide residues.

The T rich nucleic acid may include one or more CpG motifs. The motifs may be methylated or unmethylated. In other embodiments the T rich nucleic acid is free of one or more CpG dinucleotides.

In other embodiments the T rich nucleic acid has poly A, poly G, and/or poly C motifs. In other embodiments the T rich nucleic acid is free of two poly C sequences of at least 3 contiguous C nucleotide residues. Preferably the T rich nucleic acid is free of two poly A sequences of at least 3 contiguous A nucleotide residues. In other embodiments the T rich nucleic acid comprises a nucleotide composition of greater than 25% C or greater than 25% A. In yet other embodiments the T rich nucleic acid is free of poly-C sequences, poly G sequences or poly-A sequences.

In some cases the T rich nucleic acid may be free of poly T motifs, but rather, comprises a nucleotide composition of greater than 25% T. In other embodiments the T rich nucleic acid may have poly T motifs and also comprise a nucleotide composition of greater than 25% T. In some embodiments the T rich nucleic acid comprises a nucleotide composition of greater than 25% T, greater than 30% T, greater than 40% T, greater than 50% T, greater than 60% T, greater than 80% T, or greater than 90% T nucleotide residues. . The T rich nucleic acid in some embodiments is selected from the group consisting of SEQ ID NOs: 59-63, 73-75, 142, 215, 226, 241, 267-269, 282, 301, 304, 330, 342, 358, 370-372, 393, 433, 471, 479, 486, 491, 497, 503, 556-558, 567, 694, 793-794, 797, 833, 852, 861, 867, 868, 882, 886, 905, 907, 908, and 910-913. In other embodiments the T rich nucleic acids are sequence selected from the group consisting of SEQ ID NOs: 64, 98, 112, 146, 185, 204, 208, 214, 224, 233, 244, 246, 247, 258, 262, 263, 265, 270-273, 300, 305, 316, 317, 343, 344, 350, 352, 354, 374, 376, 392, 407, 411-413, 429-432, 434, 435, 443, 474, 475, 498-501, 518, 687, 692, 693, 804, 862, 883, 884, 888, 890, and 891.

In some embodiments the poly G nucleic acid comprises: 5' $X_1X_2GGGX_3X_4$ 3' wherein X_1 , X_2 , X_3 , and X_4 are nucleotides. In embodiments at least one of X_3 and X_4 are a G or both of X_3 and X_4 are a G. In other embodiments the poly G nucleic acid comprises the following formula: 5' GGGNNGGG 3' wherein N represents between 0 and 20 nucleotides. In yet other embodiments the poly G nucleic acid comprises the following formula: 5' GGGNNGGGNNGGG 3' wherein N represents between 0 and 20 nucleotides. The poly G nucleic acid in some embodiments is selected from the group consisting of SEQ ID NOs. : 5, 6, 73, 215, 267-269, 276, 282, 288, 297-299, 355, 359, 386, 387, 444, 476, 531, 557-559, 733, 768, 795, 796, 914-925, 928-931, 933-936, and 938. In other embodiments the poly G

nucleic acid includes a sequence selected from the group consisting of SEQ ID NOs; 67, 80-82, 141, 147, 148, 173, 178, 183, 185, 214, 224, 264, 265, 315, 329, 434, 435, 475, 519, 521-524, 526, 527, 535, 554, 565, 609, 628, 660, 661, 662, 725, 767, 825, 856, 857, 876, 892, 909, 926, 927, 932, and 937.

5 The poly G nucleic acid may include one or more CpG motifs or T-rich motifs. The CpG motifs may be methylated or unmethylated. In other embodiments the poly G nucleic acid is free of one or more CpG dinucleotides or poly-T motifs.

The nucleic acid molecules and optionally other agents may be administered by any route known in the art for delivering medicaments. The medicaments may be administered separately or together, in the same pharmaceutical formulation or separate formulations, by the same route or by different routes. In one embodiment the nucleic acid molecule(s) is administered on a routine schedule. In another embodiment the other agent(s) (e.g., antiangiogenesis agents, anticancer agents) is administered on a routine schedule.

Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention.

Brief Description of the Drawings

Figure 1 is a histogram showing the effect of a CpG nucleic acid on angiogenesis as measured by hemoglobin content.

20 The drawing is not required for enablement of the claimed invention.

Detailed Description of the Invention

25 The present invention includes compositions that include antiangiogenic nucleic acids and methods of using the antiangiogenic nucleic acids for the treatment of diseases that are mediated by angiogenesis. The invention includes antiangiogenic nucleic acids having various nucleotide sequences. The present invention comprises a method of treating undesired angiogenesis in a human or animal comprising the steps of the administering to the human or animal with the undesired angiogenesis a composition comprising an effective amount of, for example, an antiangiogenic nucleic acid.

30 As used herein, the term "angiogenesis" means the generation of new blood vessels into a tissue or organ. Under normal physiological conditions, humans or animals undergo angiogenesis only in very specific restricted situations. For example, angiogenesis is

normally observed in wound healing, fetal and embryonal development and formation of the corpus luteum, endometrium and placenta. The term "endothelium" means a thin layer of flat epithelial cells that lines serous cavities, lymph vessels, and blood vessels. The term "endothelial inhibiting activity" means the capability of a molecule to inhibit angiogenesis in general and, for example, to inhibit the growth of bovine capillary endothelial cells in culture in the presence of fibroblast growth factor.

Antiangiogenic nucleic acids are effective in treating diseases or processes that are mediated by, or involve, angiogenesis. The present invention includes the method of treating an angiogenesis mediated disease with an effective amount of antiangiogenic nucleic acids. The angiogenesis mediated diseases include, but are not limited to, solid tumors; blood born tumors such as leukemias; tumor metastasis; benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; pre-malignant tumors; rheumatoid arthritis; psoriasis; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation.

Antiangiogenic nucleic acids may be useful in the treatment of disease of excessive or abnormal stimulation of endothelial cells. These diseases include, but are not limited to, intestinal adhesions, atherosclerosis, scleroderma, and hypertrophic scars, i.e., keloids. Antiangiogenic nucleic acid can be used as a birth control agent by preventing vascularization required for embryo implantation.

Antiangiogenic nucleic acids may be useful in the treatment of conditions characterized by abnormal epithelial cell proliferation, such as proliferative dermatologic disorders. These include conditions such as keloids, seborrheic keratosis, papilloma virus infection (e.g. producing verruca vulbaris, verruca plantaris, verruca plana, condylomata, etc.) and eczema.

Antiangiogenic nucleic acids may be useful in the treatment of precancerous lesions such as epithelial precancerous lesions. An epithelial precancerous lesion is a lesion of epithelial cell origin that has a propensity to develop into a cancerous condition. An example is a precancerous skin lesion. Epithelial precancerous skin lesions also arise from other proliferative skin disorders such as hemangiomas, keloids, eczema and papilloma virus infections producing verruca vulbaris, verruca plantaris and verruca planar. The symptoms of

the epithelial precancerous lesions include skin-colored or red-brown macule or papule with dry adherent scales. Actinic keratosis is the most common epithelial precancerous lesion among fair skinned individuals. It is usually present as lesions on the skin which may or may not be visually detectable. The size and shape of the lesions varies. It is a photosensitive disorder and may be aggravated by exposure to sunlight. Bowenoid actinic keratosis is another form of an epithelial precancerous lesion. In some cases, the lesions may develop into an invasive form of squamous cell carcinoma and may pose a significant threat of metastasis. Other types of epithelial precancerous lesions include hypertrophic actinic keratosis, arsenical keratosis, hydrocarbon keratosis, thermal keratosis, radiation keratosis, viral keratosis, Bowen's disease, erythroplaquia of queyrat, oral erythroplaquia, leukoplakia, and intraepidermal epithelialoma.

Antiangiogenic nucleic acids may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with antiangiogenic nucleic acids and then antiangiogenic nucleic acids may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize any residual primary tumor. In some instances it may be preferable to administer the antiangiogenic nucleic acids specifically to a site likely to harbor a metastatic lesion (that may or may not be clinically discernible at the time). A sustained release formulation implanted specifically at the site (or the tissue) where the metastatic lesion is likely to be would be suitable in these latter instances.

In some embodiments, the antiangiogenic nucleic acids of the invention do not interfere with specific receptor-ligand interactions at the cell surface of a cell, thereby causing the stimulation or inhibition of signaling through such receptors. These interactions include those involving heparin binding receptor, VEGF receptor, or EGF receptor.

In still other embodiments, the antiangiogenic nucleic acids are not antisense nucleic acids, meaning that they do not function by binding to complementary genomic DNA or RNA species within a cell and thereby inhibiting the function of said genomic DNA or RNA species. In important embodiments, the antiangiogenesis nucleic acid does not comprise a nucleic acid sequence that corresponds to a VEGF encoding sequence (or is complementary to a VEGF encoding sequence).

The effective dosage for inhibition of angiogenesis *in vivo*, which can be defined as inhibition of capillary endothelial cell proliferation and/or migration and/or blood vessel ingrowth, can be extrapolated from *in vitro* inhibition assays. *In vitro* assays have been

developed to screen for inhibition of angiogenesis. Events that can be tested to assess angiogenesis inhibitors include proteolytic degradation of extracellular matrix and/or basement membrane, proliferation of endothelial cells, migration of endothelial cells, and capillary tube formation. The chick chorioallantoic membrane assay (CAM), described by Taylor and Folkman (*Nature* 297:307-312, 1982), can be used to determine whether the compound is capable of inhibiting neovascularization *in vivo*.

In some embodiments, the antiangiogenic nucleic acids are administered in doses, routes and schedules (and also in therapeutic cocktails) that would not result in the stimulation of an immune response.

The effective dosage is dependent not only on the sequence of the nucleic acid molecules used for inhibition of angiogenesis, but also on the method and means of delivery, which can be localized or systemic. For example, in some applications, as in the treatment of psoriasis or diabetic retinopathy, the inhibitor preferably is delivered in a topical or ophthalmic carrier. In other applications, as in the treatment of solid tumors, the inhibitor preferably is delivered by means of a biodegradable, polymeric implant.

An "antiangiogenic nucleic acid" as used herein is any nucleic acid containing an antiangiogenic motif or backbone that inhibits capillary endothelial cell proliferation and/or migration and/or blood vessel ingrowth.

The compounds useful according to the invention are nucleic acids. The nucleic acids may be double-stranded or single-stranded. Generally, double-stranded molecules may be more stable *in vivo*, while single-stranded molecules may have increased activity. The terms "nucleic acid" and "oligonucleotide" refer to multiple nucleotides (i.e. molecules comprising a sugar (e.g. ribose or deoxyribose) linked to a phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g. cytosine (C), thymine (T) or uracil (U)) or a substituted purine (e.g. adenine (A) or guanine (G)) or a modified base. As used herein, the terms refer to oligoribonucleotides as well as oligodeoxyribonucleotides. The terms shall also include polynucleosides (i.e. a polynucleotide minus the phosphate) and any other organic base containing polymer. Nucleic acid molecules as used herein include vectors, e.g., plasmids, as well as oligonucleotides.

The terms "nucleic acid" and "oligonucleotide" also encompass nucleic acids or oligonucleotides with a covalently modified base and/or sugar. For example, they include nucleic acids having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group

at the 5' position. Thus modified nucleic acids may include a 2'-O-alkylated ribose group. In addition, modified nucleic acids may include sugars such as arabinose instead of ribose. Thus the nucleic acids may be heterogeneous in backbone composition thereby containing any possible combination of polymer units linked together such as peptide- nucleic acids (which have amino acid backbone with nucleic acid bases). In some embodiments the nucleic acids are homogeneous in backbone composition.

The substituted purines and pyrimidines of the nucleic acids include standard purines and pyrimidines such as cytosine as well as base analogs such as C-5 propyne substituted bases (Wagner et al., *Nature Biotechnology* 14:840- 844, 1996). Purines and pyrimidines include but are not limited to adenine, cytosine, guanine, thymine, 5-methylcytosine, 2-aminopurine, 2-amino-6-chloropurine, 2,6-diaminopurine, hypoxanthine, and other naturally and non-naturally occurring nucleobases, substituted and unsubstituted aromatic moieties.

The nucleic acid is a linked polymer of bases or nucleotides. As used herein with respect to linked units of a nucleic acid, "linked" or "linkage" means two entities are bound to one another by any physicochemical means. Any linkage known to those of ordinary skill in the art, covalent or non-covalent, is embraced. Such linkages are well known to those of ordinary skill in the art. Natural linkages, which are those ordinarily found in nature connecting the individual units of a nucleic acid, are most common. The individual units of a nucleic acid may be linked, however, by synthetic or modified linkages.

Whenever a nucleic acid is represented by a sequence of letters it will be understood that the nucleotides are in 5'→ 3' order from left to right and that "A" denotes adenosine, "C" denotes cytosine, "G" denotes guanosine, "T" denotes thymidine, and "U" denotes uracil unless otherwise noted.

Nucleic acid molecules useful according to the invention can be obtained from natural nucleic acid sources (e.g. genomic nuclear or mitochondrial DNA or cDNA), or are synthetic (e.g. produced by oligonucleotide synthesis). Nucleic acids isolated from existing nucleic acid sources are referred to herein as native, natural, or isolated nucleic acids. The nucleic acids useful according to the invention may be isolated from any source, including eukaryotic sources, prokaryotic sources, nuclear DNA, mitochondrial DNA, etc. Thus, the term nucleic acid encompasses both synthetic and isolated nucleic acids.

The term "isolated" as used herein refers to a nucleic acid which is substantially free of or which is separated from components which it is normally associated with in nature e.g.,

nucleic acids, proteins, lipids, carbohydrates or *in vivo* systems to an extent practical and appropriate for its intended use. In particular, the nucleic acids are sufficiently pure and are sufficiently free from other biological constituents of host cells so as to be useful in, for example, producing pharmaceutical preparations. Because an isolated nucleic acid of the invention may be admixed with a pharmaceutically-acceptable carrier in a pharmaceutical preparation, the nucleic acid may comprise only a small percentage by weight of the preparation. The nucleic acid is nonetheless substantially pure in that it has been substantially separated from the substances with which it may be associated in living systems. The nucleic acids can be produced on a large scale in plasmids, (see Sambrook, T., *et al.*, "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor laboratory Press, New York, 1989) and separated into smaller pieces or administered whole. After being administered to a subject the plasmid can be degraded into oligonucleotides. One skilled in the art can purify viral, bacterial, eukaryotic, etc. nucleic acids using standard techniques, such as those employing restriction enzymes, exonucleases or endonucleases.

For use in the instant invention, the nucleic acids can be synthesized *de novo* using any of a number of procedures well known in the art. For example, the b-cyanoethyl phosphoramidite method (Beaucage, S.L., and Caruthers, M.H., *Tet. Let.* 22:1859, 1981); nucleoside H-phosphonate method (Garegg *et al.*, *Tet. Let.* 27:4051-4054, 1986; Froehler *et al.*, *Nucl. Acid. Res.* 14:5399-5407, 1986, ; Garegg *et al.*, *Tet. Let.* 27:4055-4058, 1986, Gaffney *et al.*, *Tet. Let.* 29:2619-2622, 1988). These chemistries can be performed by a variety of automated oligonucleotide synthesizers available in the market.

In some embodiments, the nucleic acids useful according to the invention may function as immunostimulatory nucleic acids. An immunostimulatory nucleic acid is any nucleic acid, as described herein, which is capable of modulating an immune response. A nucleic acid which modulates an immune response is one which produces any form of immune stimulation, including, but not limited to, induction of a cytokine, B cell activation, T cell activation, monocyte activation. Immunostimulatory nucleic acids include, but are not limited to, CpG nucleic acids, T-rich nucleic acids, poly G nucleic acids, and nucleic acids having phosphate modified backbones, such as phosphorothioate backbones.

A "CpG nucleic acid" or a "CpG antiangiogenic nucleic acid" as used herein is a nucleic acid containing at least one unmethylated CpG dinucleotide (cytosine-guanine dinucleotide sequence, i.e. "CpG DNA" or DNA containing a 5' cytosine followed by 3' guanosine and linked by a phosphate bond) and inhibits angiogenesis. The entire CpG nucleic


acid can be unmethylated or portions may be unmethylated but at least the C of the 5' CG 3' must be unmethylated.

In one embodiment the invention provides a CpG nucleic acid represented by at least the formula:



wherein X_1 and X_2 are nucleotides and N is any nucleotide and N_1 and N_2 are nucleic acid sequences composed of from about 0-25 N 's each. In some embodiments X_1 is adenine, guanine, or thymine and/or X_2 is cytosine, adenine, or thymine. In other embodiments X_1 is cytosine and/or X_2 is guanine.

In other embodiments the CpG nucleic acid is represented by at least the formula:



wherein X_1 , X_2 , X_3 , and X_4 are nucleotides. In some embodiments, X_1X_2 are nucleotides selected from the group consisting of: GpT, GpG, GpA, ApA, ApT, ApG, CpT, CpA, CpG, TpA, TpT, and TpG; and X_3X_4 are nucleotides selected from the group consisting of: TpT, CpT, ApT, TpG, ApG, CpG, TpC, ApC, CpC, TpA, ApA, and CpA; N is any nucleotide and N_1 and N_2 are nucleic acid sequences composed of from about 0-25 N 's each. In some embodiments, X_1X_2 are GpA or GpT and X_3X_4 are TpT. In other embodiments X_1 or X_2 or both are purines and X_3 or X_4 or both are pyrimidines or X_1X_2 are GpA and X_3 or X_4 or both are pyrimidines.

In another embodiment the CpG nucleic acid has the sequence
 $5'TCN_1TX_1X_2CGX_3X_43'$.

Examples of CpG nucleic acids according to the invention include but are not limited to those listed in Table 1, such as SEQ ID NOs: 1, 3, 4, 14-16, 18-24, 28, 29, 33-46, 49, 50, 52-56, 58, 64-67, 69, 71, 72, 76-87, 90, 91, 93, 94, 96, 98, 102-124, 126-128, 131-133, 136-141, 146-150, 152-153, 155-171, 173-178, 180-186, 188-198, 201, 203-214, 216-220, 223, 224, 227-240, 242-256, 258, 260-265, 270-273, 275, 277-281, 286-287, 292, 295-296, 300, 302, 305-307, 309-312, 314-317, 320-327, 329, 335, 337-341, 343-352, 354, 357, 361-365, 367-369, 373-376, 378-385, 388-392, 394, 395, 399, 401-404, 406-426, 429-433, 434-437, 439, 441-443, 445, 447, 448, 450, 453-456, 460-464, 466-469, 472-475, 477, 478, 480, 483-485, 488, 489, 492, 493, 495-502, 504-505, 507-509, 511, 513-529, 532-541, 543-555, 564-566, 568-576, 578, 580, 599, 601-605, 607-611, 613-615, 617, 619-622, 625-646, 648-650, 653-664, 666-697, 699-706, 708, 709, 711-716, 718-732, 736, 737, 739-744, 746, 747, 749-761, 763, 766-767, 769, 772-779, 781-783, 785-786, 7900792, 798-799, 804-808, 810, 815,

817, 818, 820-832, 835-846, 849-850, 855-859, 862, 865, 872, 874-877, 879-881, 883-885, 888-904, and 909-913.

A "T rich nucleic acid" or "T rich antiangiogenic nucleic acid" is a nucleic acid which includes at least one poly T sequence and/or which has a nucleotide composition of greater than 25% T nucleotide residues and which inhibits angiogenesis. A nucleic acid having a poly-T sequence includes at least four Ts in a row, such as 5'-TTTT-3'. Preferably the T rich nucleic acid includes more than one poly T sequence. In preferred embodiments the T rich nucleic acid may have 2, 3, 4, etc poly T sequences, such as SEQ ID NO:246 or SEQ ID NO:433. Other T rich nucleic acids have a nucleotide composition of greater than 25% T nucleotide residues, but do not necessarily include a poly T sequence. In these T rich nucleic acids the T nucleotide residues may be separated from one another by other types of nucleotide residues, i.e., G, C, and A. In some embodiments the T rich nucleic acids have a nucleotide composition of greater than 30%, 40%, 50%, 60%, 70%, 80%, 90%, and 99%, T nucleotide residues and every integer % in between. Preferably the T rich nucleic acids have at least one poly T sequence and a nucleotide composition of greater than 25% T nucleotide residues.

In one embodiment the T rich nucleic acid is represented by at least the formula:



wherein X_1 , X_2 , X_3 , and X_4 are nucleotides. In one embodiment X_1X_2 is TT and/or X_3X_4 is TT. In another embodiment X_1X_2 are any one of the following nucleotides TA, TG, TC, AT, AA, AG, AC, CT, CC, CA, CG, GT, GG, GA, and GC; and X_3X_4 are any one of the following nucleotides TA, TG, TC, AT, AA, AG, AC, CT, CC, CA, CG, GT, GG, GA, and GC.

In some embodiments it is preferred that the T-rich nucleic acid does not contain poly C (CCCC), poly A (AAAA), poly G (GGGG), CpG motifs, or multiple GGs. In other embodiments the T-rich nucleic acid includes these motifs. Thus in some embodiments of the invention the T rich nucleic acids include CpG dinucleotides and in other embodiments the T rich nucleic acids are free of CpG dinucleotides. The CpG dinucleotides may be methylated or unmethylated.

Examples of T rich nucleic acids that are free of CpG nucleic acids include but are not limited to those listed in Table 1, such as SEQ ID Nos: 59-63, 73-75, 142, 215, 226, 241, 267-269, 282, 301, 304, 330, 342, 358, 370-372, 393, 433, 471, 479, 486, 491, 497, 503, 556-558, 567, 694, 793-794, 797, 833, 852, 861, 867, 868, 882, 886, 905, 907, 908, and 910-913. Examples of T rich nucleic acids that include CpG nucleic acids include but are not limited to

those listed in Table 1, such as SEQ ID Nos: 64, 98, 112, 146, 185, 204, 208, 214, 224, 233, 244, 246, 247, 258, 262, 263, 265, 270-273, 300, 305, 316, 317, 343, 344, 350, 352, 354, 374, 376, 392, 407, 411-413, 429-432, 434, 435, 443, 474, 475, 498-501, 518, 687, 692, 693, 804, 862, 883, 884, 888, 890, and 891.

5 Poly G containing nucleic acids are also useful in accordance with the invention. A "poly G nucleic acid" or "poly G antiangiogenic nucleic acid" is a nucleic acid which includes at least one poly G sequence and/or which has a nucleotide composition of greater than 25% G nucleotide residues and which inhibits angiogenesis. A variety of references, including Pisetsky and Reich, 1993 *Mol. Biol. Reports*, 18:217-221; Krieger and Herz, 1994, *Ann. Rev. Biochem.*, 63:601-637; Macaya et al., 1993, *PNAS*, 90:3745-3749; Wyatt et al., 1994, *PNAS*, 91:1356-1360; Rando and Hogan, 1998, In *Applied Antisense Oligonucleotide Technology*, ed. Krieg and Stein, p. 335-352; and Kimura et al., 1994, *J. Biochem.* 116, 991-994 describe the properties of poly G nucleic acids.

Poly G nucleic acids preferably are nucleic acids having the following formulas:



wherein X_1 , X_2 , X_3 , and X_4 are nucleotides. In preferred embodiments at least one of X_3 and X_4 are a G. In other embodiments both of X_3 and X_4 are a G. In yet other embodiments the preferred formula is 5' GGGN GGG 3', or 5' GGGN GGGN GGG 3' wherein N represents between 0 and 20 nucleotides. In other embodiments the Poly G nucleic acid is free of

20 unmethylated CG dinucleotides, such as, for example, the nucleic acids listed below as SEQ ID Nos: 5, 6, 73, 215, 267-269, 276, 282, 288, 297-299, 355, 359, 386, 387, 444, 476, 531, 557-559, 733, 768, 795, 796, 914-925, 928-931, 933-936, and 938. In other embodiments the poly G nucleic acid includes at least one unmethylated CG dinucleotide, such as, for example, the nucleic acids listed above as SEQ ID Nos; 67, 80-82, 141, 147, 148, 173, 178, 183, 185, 214, 224, 264, 265, 315, 329, 434, 435, 475, 519, 521-524, 526, 527, 535, 554, 565, 609, 628, 25 660, 661, 662, 725, 767, 825, 856, 857, 876, 892, 909, 926, 927, 932, and 937.

The antiangiogenic nucleic acids of the invention can also be those which do not possess CpG, poly-G, or T-rich motifs.

Nucleic acids having modified backbones, such as phosphorothioate backbones, also 30 fall within the class of immunostimulatory nucleic acids. U.S. Patents Nos. 5,723,335 and 5,663,153 issued to Hutcherson, et al. and related PCT publication WO95/26204 describe immune stimulation using phosphorothioate oligonucleotide analogues. These patents

describe the ability of the phosphorothioate backbone to stimulate an immune response in a non-sequence specific manner.

The antiangiogenic nucleic acid molecules may be any size of at least 6 nucleotides but in some embodiments are in the range of between 6 and 100 or in some embodiments between 8 and 35 nucleotides in size. Nucleic acids can be produced on a large scale in plasmids. These may be administered in plasmid form or alternatively they can be degraded into oligonucleotides before administration.

“Palindromic sequence” shall mean an inverted repeat (i.e. a sequence such as ABCDEE'D'C'B'A' in which A and A' are bases capable of forming the usual Watson-Crick base pairs and which includes at least 6 nucleotides in the palindrome. *In vivo*, such sequences may form double-stranded structures. In one embodiment the nucleic acid contains a palindromic sequence. In some embodiments when the nucleic acid is a CpG nucleic acid, a palindromic sequence used in this context refers to a palindrome in which the CpG is part of the palindrome, and optionally is the center of the palindrome. In another embodiment the nucleic acid is free of a palindrome. A nucleic acid that is free of a palindrome does not have any regions of 6 nucleotides or greater in length which are palindromic. A nucleic acid that is free of a palindrome can include a region of less than 6 nucleotides which are palindromic.

A “stabilized nucleic acid molecule” shall mean a nucleic acid molecule that is relatively resistant to *in vivo* degradation (e.g. via an exo- or endo-nuclease). Stabilization can be a function of length or secondary structure. Nucleic acids that are tens to hundreds of kbs long are relatively resistant to *in vivo* degradation. For shorter nucleic acids, secondary structure can stabilize and increase their effect. For example, if the 3' end of an oligonucleotide has self-complementarity to an upstream region, so that it can fold back and form a sort of stem loop structure, then the oligonucleotide becomes stabilized and therefore exhibits more activity.

Some stabilized oligonucleotides of the instant invention have a modified backbone. It has been demonstrated that modification of the oligonucleotide backbone provides enhanced activity of the nucleic acids when administered *in vivo*. Nucleic acids, including at least two phosphorothioate linkages at the 5' end of the oligonucleotide and multiple phosphorothioate linkages at the 3' end, preferably 5, may provide maximal activity and protect the oligonucleotide from degradation by intracellular exo- and endo-nucleases. Other modified oligonucleotides include phosphodiester modified oligonucleotide, combinations of phosphodiester and phosphorothioate oligonucleotide, methylphosphonate,

methyolphosphorothioate, phosphorodithioate, and combinations thereof. Each of these combinations and their particular effects on immune cells is discussed in more detail in PCT Published Patent Applications claiming priority to U.S. Patent Nos. 6,207,646B1 and 6,239,116B1, the entire contents of which are hereby incorporated by reference. It is believed that these modified oligonucleotides may show more antiangiogenic activity due to enhanced nuclease resistance, increased cellular uptake, increased protein binding, and/or altered intracellular localization.

Other stabilized oligonucleotides include: nonionic DNA analogs, such as alkyl- and aryl-phosphates (in which the charged phosphonate oxygen is replaced by an alkyl or aryl group), phosphodiester and alkylphosphotriesters, in which the charged oxygen moiety is alkylated. Oligonucleotides which contain diol, such as tetraethyleneglycol or hexaethyleneglycol, at either or both termini have also been shown to be substantially resistant to nuclease degradation.

For use *in vivo*, nucleic acids are preferably relatively resistant to degradation (*e.g.*, via endo- and exo-nucleases). Secondary structures, such as stem loops, can stabilize nucleic acids against degradation. Alternatively, nucleic acid stabilization can be accomplished via phosphate backbone modifications. One type of stabilized nucleic acid has at least a partial phosphorothioate modified backbone. Phosphorothioates may be synthesized using automated techniques employing either phosphoramidate or H-phosphonate chemistries.

Aryl- and alkyl-phosphonates can be made, *e.g.*, as described in U.S. Patent No. 4,469,863; and alkylphosphotriesters (in which the charged oxygen moiety is alkylated as described in U.S. Patent No. 5,023,243 and European Patent No. 092,574) can be prepared by automated solid phase synthesis using commercially available reagents. Methods for making other DNA backbone modifications and substitutions have been described (Uhlmann, E. and Peyman, A., *Chem. Rev.* 90:544, 1990; Goodchild, J., *Bioconjugate Chem.* 1:165, 1990).

Other sources of nucleic acids useful according to the invention include standard viral and bacterial vectors, many of which are commercially available. In its broadest sense, a "vector" is any nucleic acid material which is ordinarily used to deliver and facilitate the transfer of nucleic acids to cells. The vector as used herein may be an empty vector or a vector carrying a gene which can be expressed. In the case when the vector is carrying a gene the vector generally transports the gene to the target cells with reduced degradation relative to the extent of degradation that would result in the absence of the vector. In this case the vector

optionally includes gene expression sequences to enhance expression of the gene in target cells such as immune cells, but it is not required that the gene be expressed in the cell.

In general, vectors include, but are not limited to, plasmids, phagemids, viruses, other vehicles derived from viral or bacterial sources. Viral vectors are one type of vector and include, but are not limited to, nucleic acid sequences from the following viruses: retrovirus, such as Moloney murine leukemia virus, Harvey murine sarcoma virus, murine mammary tumor virus, and Rous sarcoma virus; adenovirus, adeno-associated virus; SV40-type viruses; polyoma viruses; Epstein-Barr viruses; papilloma viruses; herpes virus; vaccinia virus; polio virus; and RNA virus such as a retrovirus. One can readily employ other vectors not named but known to the art. Some viral vectors are based on non-cytopathic eukaryotic viruses in which non-essential genes have been replaced with a nucleic acid to be delivered. Non-cytopathic viruses include retroviruses, the life cycle of which involves reverse transcription of genomic viral RNA into DNA.

Standard protocols for producing empty vectors or vectors carrying genes (including the steps of incorporation of exogenous genetic material into a plasmid, transfection of a packaging cell line with plasmid, production of recombinant retroviruses by the packaging cell line, collection of viral particles from tissue culture media, and/or infection of the target cells with viral particles) are provided in Kriegler, M., "Gene Transfer and Expression, A Laboratory Manual," W.H. Freeman C.O., New York (1990) and Murry, E.J. Ed. "Methods in Molecular Biology," vol. 7, Humana Press, Inc., Clifton, New Jersey (1991).

Other vectors include plasmid vectors. Plasmid vectors have been extensively described in the art and are well-known to those of skill in the art. See e.g., Sambrook et al., "Molecular Cloning: A Laboratory Manual," Second Edition, Cold Spring Harbor Laboratory Press, 1989. In the last few years, plasmid vectors have been found to be particularly advantageous for delivering genes to cells *in vivo* because of their inability to replicate within and integrate into a host genome. Some plasmids, however, having a promoter compatible with the host cell, can express a peptide from a gene operatively encoded within the plasmid. Some commonly used plasmids include pBR322, pUC18, pUC19, pcDNA3.1, SV40, and pBlueScript. Other plasmids are well-known to those of ordinary skill in the art. Additionally, plasmids may be custom designed using restriction enzymes and ligation reactions to remove and add specific fragments of DNA.

Exemplary antiangiogenic nucleic acid sequences include but are not limited to those antiangiogenic sequences shown in Table 1 (SEQ ID NO: 1 to SEQ ID NO:1093). The Table

lists the SEQ ID NO, nucleotide sequence of the oligonucleotide (ODN sequence), and backbone modification, if any.

Backbone modifications are abbreviated as follows:

S = phosphorothioate

O = phosphodiester

SOS = phosphorothioate and phosphodiester chimeric with phosphodiester in middle

SO = phosphorothioate and phosphodiester chimeric with phosphodiester on 3' end

OS = phosphorothioate and phosphodiester chimeric with phosphodiester on 5' end

S2 = phosphorodithioate

S2O = phosphorodithioate and phosphodiester chimeric with phosphodiester on 3' end

OS2 = phosphorodithioate and phosphodiester chimeric with phosphodiester on 5' end

X = unknown

p-ethoxy = p-ethoxy backbone; see, e.g., US patent 6,015,886

PO = phosphodiester

ODN sequence symbols, other than a, c, g and t, are as follows:

i = inosine

n = a, c, g, or t

d = a, g or t

h = a, c or t

b = c, g or t; if "b" is single and is listed on 5' or 3' end of oligonucleotide, then "b" indicates a biotin moiety attached to that end of the oligonucleotide

q = 5-methyl-cytosine

m = a or c

s = c or g

x = if "x" is single and is listed on 5' or 3' end of oligonucleotide, then "x" indicates a biotin moiety attached to that end of the oligonucleotide

z = 5-methyl-cytidine

f = FITC moiety attached to 5' or 3' end of oligonucleotide

Table 1

SEQ ID NO:	ODN SEQUENCE	BACKBONE
1	tctcccagcgtgcgccat	S
2	ataatccagcttgaaccaag	S
3	ataatcgacgttcaagcaag	S
4	taccgcgtgcgaccctct	S
5	ggggagggt	S

TABLE "E6/100"

6	ggggagggg	S
7	ggtgaggtg	S
8	tccatgtzgttcctgatgct	O
9	gctaccttagzgtga	O
10	tccatgazgttcctgatgct	O
11	tccatgacgttcztgatgct	O
12	gctagazgttagtgt	S
13	agctccatggtgctcactg	S
14	ccacgtcgaccctcaggcga	S
15	gcacatcgtcccgcagccga	S
16	gtcactcgtggtacctcga	O
17	gttggatacaggccagactttgttg	O
18	gattcaacttgcgctcatcttaggc	S
19	accatggacgaactgtttcccctc	S
20	accatggacgagctgtttcccctc	S
21	accatggacgacctgtttcccctc	S
22	accatggacgtactgtttcccctc	S
23	accatggacggtctgtttcccctc	S
24	accatggacgttctgtttcccctc	O
25	ccactcacatctgctgctccacaag	O
26	acttctcatagtccctttggtccag	O
27	tccatgagcttcctgagtct	O
28	gaggaaggigiggaigacgt	O
29	gtgaaticgttcicgggict	S
30	aaaaaa	S
31	cccccc	S
32	ctgtca	S
33	tcgtag	S
34	tcgtgg	S
35	cgtcgt	SOS
36	tccatgtcggtcctgagtct	SOS
37	tccatgccggtcctgagtct	SOS
38	tccatgacggtcctgagtct	SOS
39	tccatgacggtcctgagtct	SOS
40	tccatgtcgatcctgagtct	SOS
41	tccatgtcgctcctgagtct	SOS
42	tccatgtcggtcctgagtct	SOS
43	tccatgacgttcctgagtct	SOS
44	tccataacgttcctgagtct	SOS
45	tccatgacgtccctgagtct	SOS
46	tccatcacgtgctgagtct	SOS
47	tccatgctggtcctgagtct	SOS
48	tccatgtzgttcctgagtct	O
49	ccgcttcctccagatgagctcatgggtttctccaccaag	O
50	cttgggtggagaaacccatgagctcatctggaggaagcgg	O
51	cccaaagggatgagaagtt	O
52	agatagcaaatacggtgacg	O
53	ggttcacgtgctcatggctg	S
54	tctcccagcgtgcgccat	S
55	tctcccagcgtgcgccat	S
56	taccgctgcgaccctct	S
57	ataatccagcttgaaccaag	S
58	ataatcgacgttcaagcaag	O
59	tccatgattttcctgatttt	S
60	ttgtttttttgtttttttgttttt	O
61	ttttttttgtttttttgttttt	S
62	tgctgcttttgtgcttttgtgctt	S

63	tgctgcttgctgttttgctgctt	O
64	gcattcatcaggcgggcaagaat	O
65	taccgagcttcgacgagatttca	O
66	gcatgacgttgagct	S
67	cacgttgaggggcat	S
68	ctgctgagactggag	S
69	tccatgacgttcctgacgtt	O
70	gcatgagcttgagctga	S
71	tcagcgtgcgcc	S
72	atgacgttcctgacgtt	S
73	ttttgggggttttgggggtttt	S
74	tctaggcttttttaggcttcc	S
75	tgcatttttttaggccaccat	S
76	tctcccagcgtgcgtgcgccat	S
77	tctcccagcgggcgcac	S
78	tctcccagcgcgcgccat	S
79	tctcccagcgcgcgccat	SOS
80	gggggtgacgttcagggggg	SOS
81	gggggtccagcgtgcgccatggggg	SOS
82	gggggtgcgttcagggggg	S
83	tccatgtcgttcctgtcgtt	S
84	tccatagcgttcctagcgtt	S
85	tcgtcgtgtctccgcttctt	SOS
86	gcatgacgttgagct	SOS
87	tctcccagcgtgcgccatat	S
88	tccatgazgttcctgazgtt	O
89	gcatgazgttgagct	SOS
90	tccagcgtgcgccata	O
91	tctcccagcgtgcgccat	O
92	tccatgagcttcctgagctt	SOS
93	gcatgtcgttgagct	S
94	tcctgacgttcctgacgtt	O
95	gcatgatgttgagct	O
96	gcatttcgaggagct	O
97	gcatgtagctgagct	O
98	tccaggacgttcctagttct	O
99	tccaggagcttcctagttct	O
100	tccaggatgttcctagttct	O
101	tccagtctaggcctagttct	O
102	tccagttcgagcctagttct	SOS
103	gcatggcgttgagct	SOS
104	gcatagcgttgagct	SOS
105	gcattgcgttgagct	SOS
106	gcttgcggttgcggtt	SOS
107	tctcccagcgttgcgccatat	SOS
108	tctcccagcgtgcgttatat	SOS
109	tctccctgcgtgcgccatat	SOS
110	tctgcgtgcgtgcgccatat	SOS
111	tctcctagcgtgcgccatat	SOS
112	tctcccagcgtgcgcctttt	O
113	gctandcghhagc	O
114	tcctgacgttccc	O
115	ggaagacgttaga	O
116	tcctgacgttaga	O
117	tcagaccagctggtcgggtgttcctga	O
118	tcaggaacacccgaccagctggctga	O
119	gctagtcgatagc	O

1001661001

001995-1001

120	gctagtcgctagc	o
121	gcttgacgtctagc	o
122	gcttgacgttttagc	o
123	gcttgacgtcaagc	o
124	gctagacgttttagc	o
125	tccatgacattcctgatgct	o
126	gctagacgtctagc	o
127	ggctatgtcgttcctagcc	o
128	ggctatgtcgatcctagcc	o
129	ctcatgggtttctccaccaag	o
130	cttgggtggagaaacccatgag	o
131	tccatgacgttcctagttct	o
132	ccgcttcctccagatgagctcatg	o
133	catgagctcatctggaggaagcgg	o
134	ccagatgagctcatgggtttctcc	o
135	ggagaaacccatgagctcatctgg	o
136	agcatcaggaacgacatgga	RNA
137	tccatgacgttcctgacgtt	o
138	gcgcgcgcgcgcgcgcgcgcg	o
139	ccggccggccggccggccggccgg	o
140	ttccaatcagccccaccgcgtctggccccaccctcaccctcca	o
141	tggagggtgaggggtggggccagagcgggtgggggtgattggaa	o
142	tcaaagtgtgggattttcccatgagctt	o
143	agactcatgggaaaatcccacatttga	o
144	tgccaagtgtgagtcactaataaaga	o
145	tctttattagtgactcagcacttggca	o
146	tgcaggaagtccgggttttccccaacccccc	o
147	gggggggttggggaaaacccggacttcctgca	Sos
148	ggggactttccgctggggactttccagggggactttcc	o
149	tccatgacgttcctctccatgacgttcctctccatgacgttcctc	o
150	gaggaacgtcatggagaggaacgtcatggagaggaacgtcatgga	s
151	ataatagagcttcaagcaag	s
152	tccatgacgttcctgacgtt	sos
153	tccatgacgttcctgacgtt	s
154	tccaggactttcctcaggtt	o
155	tcttgcatgctaaaggacgtcacattgcacaatcttaataaggt	o
156	accttattaagattgtgcaatgtgacgtcctttagcatcgcaaga	o
157	tcttgacgttcctggcgggtcctgtcgct	o
158	tctgtcgctcctgtcgct	o
159	tctgacgttgaagt	o
160	tctgtcggttgaagt	o
161	tctggcggttgaagt	o
162	tctgccgttgaagt	o
163	tcttacgttgaagt	o
164	tctaacgttgaagt	o
165	tctcacgttgaagt	o
166	tctgacgatgaagt	o
167	tctgacgtgaagt	o
168	tctgacggtgaagt	o
169	tctgacgtagaagt	o
170	tctgacgtcgaagt	o
171	tctgacgtggaagt	o
172	tctgagcttgaagt	o
173	gggggacgttggggg	o
174	tctgacgttccttc	s
175	tctcccagcgcgcgcgcgcgc	0
176	tctgacgttcccctggcgggtcccctgtcgct	

1001567001

177	tctgtcgctcctgtcgctcctgtcgct	0
178	tctggcggggaagt	o
179	tctgazgttgaagt	o
180	tcztgacgttgaagt	o
181	tcctagcgttgaagt	o
182	tccagacgttgaagt	o
183	tcctgacggggaagt	o
184	tcctggcggggaagt	o
185	ggctccggggagggaatttttgtctat	o
186	atagacaaaaattccctccccggagcc	RNA
187	tccatgagcttccttgagtct	so
188	tcgtcgctgtctccgcttctt	s20
189	tcgtcgctgtctccgcttctt	0
190	tcgagacattgcacaatcatctg	0
191	cagattgtgcaatgtctcga	0
192	tccatgtcgcttcctgatgcg	0
193	gcgatgtcgcttcctgatgct	0
194	gcgatgtcgcttcctgatgcg	0
195	tccatgtcgcttcgcgcgcgcg	0
196	tccatgtcgcttcctgcccgt	0
197	tccatgtcgcttcctgtagct	0
198	gcggcggggcggcgcgcgcgc	0
199	atcaggaacgtcatgggaagc	p-ethoxy
200	tccatgagcttcctgagtct	p-ethoxy
201	tcaacgtt	p-ethoxy
202	tcaagctt	S
203	tctgtcgcttcctgtcggt	S
204	tccatgtcgctttttgtcggt	S
205	tctgtcgcttcctgtcggt	S
206	tcttgtcgcttcctgtcggt	os
207	btccattccatgacgttcctgatgcttcca	S
208	tctgtcgctttttgtcggt	S
209	tcgtcgctgtctccgcttctt	s
210	tcgtcgctgtctgcccttctt	s
211	tcgtcgctgttgcgtttctt	o
212	tctgtcgcttcctgtcggttggaacgacagg	o
213	tctgtcgcttcctgtcggttcaacgtcaggaacgacagga	sos
214	ggggtctgtcgcttttgggggg	sos
215	ggggtctgtcgcttttgggggg	o
216	tccggccggtgaagt	o
217	tccggacggtgaagt	o
218	tcccgccggtgaagt	o
219	tccagacggtgaagt	o
220	tcccgacggtgaagt	o
221	tccagagcttgaagt	s
222	tccatgtzgttcctgtzgtt	sos
223	tccatgacgttcctgacgtt	sos
224	ggggttgacgttttgggggg	s
225	tccaggacttcttcaggtt	s
226	tttttttttttttttttttt	s
227	tccatgccgttcctgccgtt	s
228	tccatggcgggcctggcggg	s
229	tccatgacgttcctgccgtt	s
230	tccatgacgttcctggcggg	s
231	tccatgacgttcctgcgttt	s
232	tccatgacggtcctgacggt	s
233	tccatgcgtgcgtgcgtttt	s

234	tccatgcgttgcggtgcgtt	s
235	Btccattccattcttaggcctgagttcccat	os
236	tccatagcggtcctagcggt	o
237	tccatgtcggtcctgtcggt	o
238	tccatagcgatcctagcgat	o
239	tccattgcgttccttgcggt	o
240	tccatagcggtcctagcggt	
241	tccatgattttcctgcagttcctgatttt	s
242	tccatgacgttcctgcagttcctgacgtt	O
243	ggcggcggcggcggcggcggcg	S
244	tccacgacgttttcgacgtt	S
245	tcgtcggtgtcggtgtcggt	S
246	tcgtcggtttgtcggtttgtcggt	S
247	tcgtcggtgtcggtttgtcggt	S
248	gcgtgcgttgctggtgtcggt	o
249	czggczggczggczggczgg	S
250	gcggcgggcggcgcgcgcgc	o
251	agicccgigaacgiattcac	S
252	tgctggttgctggttgctggt	S
253	tgctggtgtcggtgtcggtgtcggt	S
254	tgctggtgtcggtgtcggtgtcggt	s
255	tcgtcgctcgctggt	s
256	tgctggtgtcggt	s
257	cccccccccccccccccccc	sos
258	tctagcggttttagcggttc	s
259	tgcacccccaggccacat	sos
260	tcgtcgctcgctcgctcgctcggt	sos
261	tcgtcggtgtcggtgtcggt	sos
262	tcgtcggtttgtcggtttgtcggt	sos
263	tcgtcggtgtcggtttgtcggt	o
264	ggggagggaggaacttcttaaaattcccccagaatgttt	o
265	aaacattctgggggaattttaagaagttcctccctcccc	o
266	atgtttacttcttaaaattcccccagaatgttt	o
267	aaacattctgggggaattttaagaagtaaacat	o
268	atgtttactagacaaaattcccccagaatgttt	o
269	aaacattctgggggaattttgtctagtaaacat	sos
270	aaaattgacgttttaaaaaa	sos
271	ccccttgacgttttcccccc	
272	ttttcggtgtttttgtcggt	sos
273	tcgtcggtttgtcggtttgtcggt	o
274	ctgcagcctgggac	o
275	accgctcgtaattatagtaaaacc	o
276	ggtacctgtggggacattgtg	o
277	agcaccgaacgtgagagg	o
278	tccatgccgttcctgccgtt	o
279	tccatgacggtcctgacggt	o
280	tccatgccggtcctgccggt	o
281	tccatgcgcgtcctgcgcgt	s
282	ctggtctttctgggtttttctgg	sos
283	tcaggggtggggggaacctt	o
284	tccatgazgttcctagtctt	o
285	tccatgatgttcctagtctt	o
286	cccgaagtcatttcctcttaacctgg	o
287	ccagggttaagaggaaatgacttcggg	o
288	tcctggzggggaagt	x
289	gzggzgggzggzgzgzcgc	o
290	tccatgtgcttcctgatgct	

291	tccatgtccttcctgatgct	
292	tccatgtcggttcctagttct	o
293	tccaagtagttcctagttct	o
294	tccatgtagttcctagttct	s
295	tcccgcgcgttcgcgcgtt	s
296	tcctggcggtcctggcgtt	o
297	tcctggaggggaagt	o
298	tcctgggggggaagt	o
299	tcctgggtgggaagt	o
300	tcgtcgttttgtcgttttgtcggt	o
301	ctgggtctttctgggtttttctgg	o
302	tccatgacgttcctgacgtt	sos
303	tccaggacttctctcaggtt	o
304	tzgtzgttttgtzgttttgtzgtt	os
305	btcgtcgttttgtcgttttgtcggtttttt	s
306	gctatgacgttccaagg	s
307	tcaacgtt	o
308	tccaggacttctctcaggtt	s
309	ctctctgtaggcccgcttgg	s
310	ctttccgttggacccctggg	s
311	gtccgggcccaggccaaagtc	s
312	gtgcgcgcgagcccgaatc	s
313	tccatgaigttcctgaigtt	o
314	aatagtcgccataacaaaaac	o
315	aatagtcgccatggcggggc	os
316	btttttccatgtcggttcctgatgcttttt	o
317	tcctgtcggttgaagttttt	o
318	gctagcttttagagcttttagagctt	o
319	tgtgcttccccccccccc	o
320	tgcacgttcccccccccccc	o
321	tcgtcggtcccccccccccc	o
322	tcgtcggtcccccccccccc	o
323	tcgccgttcccccccccccc	o
324	tcgtcgatcccccccccccc	s
325	tcctgacgttgaagt	s
326	tcctgccgttgaagt	s
327	tcctgacggtgaagt	s
328	tcctgagcttgaagt	s
329	tcctggcggggaagt	sos
330	aaaatctgtgcttttaaaaaa	o
331	gatccagtcacagtgcctggcagaatctggat	o
332	gatccagattctgccaggtcactgtgactggat	o
333	gatccagtcacagtgcctcagcagaatctggat	o
334	gatccagattctgctgagtcactgtgactggat	o
335	tcgtcggtccccccczzccc	o
336	tzgtggttcccccccccccc	o
337	tzgtcggtcccccccccccc	o
338	tcgtzgttcccccccccccc	o
339	tcgtcggtcccccccccccc	o
340	tcgtcggtcccccccccccc	o
341	tcggcggtcccccccccccc	o
342	ggccttttcccccccccccc	s
343	tcgtcgttttgacgtttttgtcggt	s
344	tcgtcgttttgacgtttttgacgtt	o
345	ccgtcggtcccccccccccc	o
346	gcgtcggtcccccccccccc	o
347	tcgtcattcccccccccccc	o

100176612301

348	acgtcggtccccccccccc	o
349	ctgtcggtccccccccccc	o
350	btttttcggtcggtccccccccccc	os
351	tcgtcggtccccccccccc	o
352	tcgtcggtttgtcggtttgtcggtb	o
353	tccagttccttcctcagtt	o
354	tzgtcggtttgtcggtttgtcggt	s
355	tcctggaggggaagt	s
356	tcctgaaaaggaagt	s
357	tcgtcggtccccccccccc	s
358	tzgtzgtttgtzgtttgtzgtt	sos
359	ggggtcaagcttgagggggg	s
360	tgctgcttccccccccccc	s2
361	tcgtcggtcggtcggt	s20
362	tcgtcggtcggtcggt	os2
363	tcgtcggtcggtcggt	s
364	tcaacgttga	s
365	tcaacgtt	
366	atagttttccatttttttac	o
367	aatagtcgcatcgcgcgac	o
368	aatagtcgcatcccgggac	o
369	aatagtcgcatcccccccc	o
370	tgctgctttgtgctttgtgctt	s
371	ctgtgctttctgtgtttctgtg	s
372	ctaatttttctaattttttctaa	s
373	tcgtcggtggtgctggtgctggt	s
374	tcgtcggtggtgctggtggtggt	
375	accatggacgagctgttccctc	s
376	tcgtcggtttgctgctggtt	
377	ctgtaagtgaagcttgagag	
378	gagaacgctggaccttcc	
379	cgggcgactcagttctatcgg	
380	gttctcagataaagcggaaccagcaacagacacagaa	
381	ttctgtgtctgttgctggttccgctttatctgagaac	
382	cagacacagaagcccgatagacg	
383	agacagacacgaaacgaccg	
384	gtctgtcccatgatctcgaa	
385	gctggccagcttacctccc	
386	ggggcctctatacaacctggg	
387	ggggtccctgagactgcc	
388	gagaacgctggaccttccat	
389	tccatgtcggtcctgatgct	
390	ctcttgcgacctggaaggta	
391	aggtacagccaggactacga	
392	accatggacgacctgttccctc	
393	accatggattaccttttccctt	o
394	atggaagggtccagcggtctc	o
395	agcatcaggaccgacatgga	
396	ctctccaagctcacttacag	
397	tccctgagactgccccacctt	
398	gccacaaaactgtccatg	
399	gtccatggcgtgcgggatga	
400	cctctatacaacctgggac	
401	cgggcgactcagttctatcgg	
402	gcgctaccggtagcctgagt	
403	cgactgccgaacaggatatcggtgatcagcactgg	
404	ccagtgtgatcaccgatatcctgttcggcagtcg	

10017995-121801

TABLE 1001

405	ccaggttgatatagaggc	s
406	tctcccagcgtacgccat	s
407	tctcccagcgtgcgtttt	s
408	tctcccagcgtgcgccat	s
409	tctcccgtcgtgcgccat	s
410	ataatcgtcgttcaagcaag	s2
411	tcgtcgttttgcgttttgcgt	s2
412	tcgtcgttttgcgttttgcgtt	s2
413	tcgtcgttttgcgttttgcgtt	s
414	tontcgtntntcgtntntcgtnt	s
415	tctcccagcgtgcgccat	s
416	tctcccatcgtgcgccat	s
417	ataatcgtgcgttcaagaaag	s
418	ataatcgacgttccccccc	s
419	tctatcgacgttcaagcaag	s
420	tcc tga cgg gg agt	
421	tccatgacgttcctgatcc	
422	tccatgacgttcctgatcc	
423	tccatgacgttcctgatcc	s
424	tcc tgg cgt gga agt	
425	tccatgacgttcctgatcc	s
426	tcgtcgtgttgcgttttctt	s
427	agcagcttttagagcttttagagctt	s
428	cccccccccccccccccccc	s
429	tcgtcgttttgcgttttgcgttttgcgtt	s
430	tcgtcgttttttgcgtttttgcgtt	s
431	tcgtcgtttttttttttttt	sos
432	tttttcaacgttgattttt	S
433	ttttttttttttttttttttt	
434	ggggtcgtcgttttgggggg	
435	tcgtcgttttgcgttttgggggg	s
436	tcgtcgtgtctccgcttcttcttgcc	s
437	tcgtcgtgtctccg	
438	ctgtaagtgcgttgagag	
439	gagaacgctggaccttccat	
440	ccaggttgatatagaggc	
441	gctagacgttagcgtga	
442	ggagctcttcgaacgccata	
443	tctccatgatggttttatcg	
444	aaggtggggcagtcctcaggga	
445	atcggaggactggcgcgccg	
446	ttaggacaaggtctagggtg	
447	accacaacgagaggaacgca	
448	ggcagtgacggctcaccggg	
449	gaaccttccatgctgtt	
450	gctagacgttagcgtga	
451	gcttggaggcctgtaagt	
452	gtagccttcccta	
453	cggtagccttcccta	
454	cacggtagccttcccta	
455	agcacggtagccttcccta	
456	gaacgctggaccttccat	
457	gaccttccat	
458	tggaccttccat	
459	gctggaccttccat	
460	acgctggaccttccat	
461	taagctctgtcaacgccagg	

1007562001

462	gagaacgctggaccttccatgt	
463	tccatgtcggtcctgatgct	
464	ttcatgccttgcaaaatggcg	
465	tgctagctgtgcctgtacct	
466	agcatcaggaccgacatgga	
467	gaccttccatgtcggtcctgat	
468	acaaccacgagaacgggaac	
469	gaaccttccatgctgttcgg	
470	caatcaatctgaggagaccc	
471	tcagctctgggtactttttca	
472	tggttacggctctgtcccatg	
473	gtctatcggaggactggcgc	
474	cattttacgggcgggcgggc	
475	gaggggaccattttacgggc	
476	tgtccagccgaggggacccat	
477	cgggcttacggcggtatgctg	
478	tggaccttctatgtcgggtcc	
479	tgtcccatgtttttagaagc	
480	gtggttacggctcgtgcccat	
481	cctccaaatgaaagaccccc	
482	ttgtactctccatgatgggt	
483	ttccatgctgttcgggtcgg	
484	gaccttctatgtcgggtcctg	
485	gagaccgctcgaccttcgat	
486	ttgccccatatttttagaaac	
487	ttgaaactgaggtgggac	
488	ctatcggaggactggcgcgcc	
489	cttggagggcctcccggcgg	
490	gctgaaccttccatgctgtt	
491	tagaaacagcattcttcttttagggcagcaca	
492	agatgggttctcagataaagcggaa	
493	ttccgctttatctgagaaccatct	
494	gtcccaggttgtatagaggctgc	
495	gcgccagtcctccgatagac	
496	atcggaggactggcgcgccg	
497	ggtctgtcccatatttttag	sos
498	tttttcaacggttgagggggg	sos
499	tttttcaagcgttgatttttt	sos
500	gggggtcaacggttgatttttt	sos
501	gggggttttcaacgttttgagggggg	
502	ggttacggctctgtcccatat	
503	ctgtcccatatttttagaca	
504	accatcctgaggccattcgg	
505	cgtctatcgggcttctgtgtctg	
506	ggccatcccacattgaaagt	
507	ccaaatatcggtgggtcaagcac	
508	gtgcttgaccaccgatatttgg	
509	gtgctgatcaccgatattcgttcgg	
510	ggccaactttcaatgtgggatggcctc	
511	ttccgcccgaatggcctcaggatggtag	
512	tatagtccctgagactgccccaccttctcaacaacc	
513	gcagcctctatacaacctgggacggga	
514	ctatcggaggactggcgcgccg	
515	tatcggaggactggcgcgccg	
516	gatcggaggactggcgcgccg	
517	ccgaacaggatatcggatgatcagcac	
518	ttttggggtcaacggttgagggggg	

519	ggggtcaacgttgagggggg	sos
520	cgcgcgcgcgcgcgcgcg	s
521	ggggcatgacgttcgggggg	ss
522	ggggcatgacgttcaaaaaa	s
523	ggggcatgagcttcgggggg	s
524	ggggcatgacgttcgggggg	sos
525	aaaacatgacgttcaaaaaa	sos
526	aaaacatgacgttcgggggg	sos
527	ggggcatgacgttcaaaaaa	sos
528	accatggacgatctgtttccctc	s
529	gccatggacgaactgttccctc	s
530	cccccccccccccccccc	sos
531	ggggggggggggggggggg	sos
532	gctgtaaaatgaatcggccg	sos
533	ttcgggcggaactcctccatt	sos
534	tatgccgcgcccggacttat	sos
535	ggggtaatcgatcagggggg	sos
536	tttgagaacgctggacctc	sos
537	gatcgctgatctaagtctcg	sos
538	gtcggtcctgatgctgttcc	sos
539	tcgtcgtcagttcgctgtcg	sos
540	ctggaccttccatgtcgg	sos
541	gctcgttcagcgctct	sos
542	ctggaccttccatgtc	sos
543	cactgtccttcgtcga	sos
544	cgctggaccttccatgtcgg	sos
545	gctgagctcatgccgtctgc	sos
546	aacgctggaccttccatgtc	sos
547	tgcattgccgtacacagctc	sos
548	ccttccatgtcggctcctgat	sos
549	tactcttcggatcccttgcg	sos
550	ttccatgtcggctcctgat	sos
551	ctgattgctctctcgtga	o
552	ggcgttattcctgactcgcc	o
553	cctacgttgtatgcgcccagct	o
554	ggggtaatcgatgagggggg	o
555	ttcgggcggaactcctccatt	o
556	ttttttttttttttttttt	o
557	gggggtttttttttttggg	o
558	tttttgggggggggttttt	o
559	gggggggggggggggggt	o
560	aaaaaaaaaaaaaaaaaaaa	o
561	cccccaaaaaaaaaaccccc	o
562	aaaaaccccccccccaaaaa	o
563	tttgaattcaggactggtgaggttgag	o
564	tttgaatcctcagcggctcctcagtgcc	o
565	aattctctatcggggcttctgtgtctgttgctggttccgctttat	o
566	ctagataaagcgggaaccagacaacagagaagccccgatagag	o
567	ttttctagagaggtgcacaatgctctgg	o
568	tttgaattccgtgtacagaagcgagaagc	o
569	tttgcgccgctagacttaacctgagagata	o
570	tttgggcccacgagagacagagacacttc	o
571	tttgggcccgttctcgttctgtacacg	s
572	gagaacgctggaccttccat	s
573	tccatgtcggctcctgatgct	s
574	ctgtcg	s
575	tcgtga	s

001551

576	cgtcga	s
577	agtgc	s
578	ctgtcg	o
579	agtgc	o
580	cgtcga	o
581	tcgtga	o
582	gagaacgctccagcttcgat	o
583	gctagacgtaagcgtga	o
584	gagaacgctcgaccttccat	o
585	gagaacgctggacctatccat	o
586	gctagaggttagcgtga	o
587	gagaacgctggacttccat	o
588	tcacgctaacgtctagc	o
589	bgctagacgcttagcgtga	o
590	atggaaggtcgagcgttctc	o
591	gagaacgctggaccttcgat	o
592	gagaacgatggaccttccat	o
593	gagaacgctggatccat	o
594	gagaacgctccagcactgat	o
595	tccatgtcggtcctgctgat	o
596	atgtcctcggtcctgatgct	o
597	gagaacgctccaccttccat	o
598	gagaacgctggaccttcgta	o
599	batggaaggtccagcgttctc	o
600	tcctga	o
601	tcaacggt	o
602	aacggt	o
603	aacggtga	o
604	tcacgctaacctctagc	o
605	gagaacgctggaccttgcat	o
606	gctggaccttccat	o
607	gagaacgctggacctcatccat	o
608	gagaacgctggacgctcatccat	o
609	aacggtgaggggcat	o
610	atgccctcaacggt	o
611	tcaacggtga	o
612	gctggaccttccat	o
613	caacggt	o
614	acaacggtga	o
615	tcacgt	o
616	tcaagcgt	o
617	tcgtca	o
618	aggatatc	o
619	tagacgtc	o
620	gacgtcat	o
621	ccatcgat	o
622	atcgatgt	o
623	atgcatgt	o
624	ccatgcat	o
625	agcgtga	o
626	tcagcgt	o
627	ccttcgat	s
628	gtgccggggtctccgggc	s
629	gctgtggggcggtcctg	o
630	btcaacggt	o
631	ftcaacggt	o
632	faacggtga	o

1007552501

633	tcaacgt	s
634	aacgttg	s
635	cgacga	o
636	tcaacgtt	o
637	tcgga	o
638	agaacgtt	o
639	tcacgat	s
640	taaacgtt	s
641	ccaacgtt	s
642	gctcga	s
643	cgacgt	s
644	cgtcgt	s
645	acgtgt	s
646	cgttcg	s
647	gagcaagctggaccttccat	s
648	cgcgta	s
649	cgtacg	s
650	tcaccggt	s
651	caagagatgctaacaatgca	s
652	acccatcaatagctctgtgc	o
653	ccatcgat	o
654	tcgacgtc	o
655	ctagcgct	o
656	taagcgct	o
657	tcgcgaattcgcg	o
658	atggaaggtccagcgttct	o
659	actggacgttagcgtga	o
660	cgccctggggctggtctgg	o
661	gtgtcggggctctccgggc	o
662	gtgccggggctctccgggc	o
663	cgccgtcgcggcggttgg	o
664	gaagttcacgttgaggggcat	s
665	atctggtgagggcaagctatg	s
666	ggtgaaacccgagaacatcat	o
667	gcaacgtt	o
668	gtaacgtt	o
669	cgaacgtt	o
670	gaaacgtt	o
671	caaacgtt	o
672	ctaacgtt	o
673	ggaacgtt	o
674	tgaacgtt	o
675	acaacgtt	o
676	ttaacgtt	o
677	aaaacgtt	o
678	ataacgtt	o
679	aacgttct	o
680	tccgatcg	o
681	tccgtacg	o
682	gctagacgctagcgtga	o
683	gagaacgctggacctcatcatccat	o
684	gagaacgctagaccttctat	o
685	actagacgttagtgtga	o
686	cacaccttgggtcaatgtcacgt	o
687	tctccatcctatgggtttatcg	o
688	cgctggaccttccat	o
689	caccaccttgggtcaatgtcacgt	o

1007995-24001

690	gctagacgcttagctgga	o
691	agtgcgattgcagatcg	o
692	ttttcgttttgtggtttgtggtt	
693	ttttcgttttgtcgttttgtcggt	
694	ttttgtttttgtggtttgtggtt	
695	accgcatggattctaggcca	s
696	gctagacgcttagcgt	o
697	aacgctggaccttccat	o
698	tcaazggt	o
699	ccttcgat	s
700	actagacgcttagtgtga	s
701	gctagaggcttagcgtga	o
702	atggactctccagcgttctc	o
703	atcgactctcgagcgttctc	o
704	gctagacgcttagc	o
705	gctagacgt	o
706	agtgcgattcgagatcg	o
707	tcagzgct	o
708	ctgattgctctctcggtga	o
709	tzaacggt	o
710	gagaazgctggaccttccat	o
711	gctagacgcttaggctga	o
712	gctacttagcgtga	o
713	gctaccttagcgtga	o
714	atcgactctcgagcgttctc	o
715	atgcactctgcagcgttctc	o
716	agtgactctccagcgttctc	o
717	gccagatgcttagcgtga	o
718	atcgactcgagcgttctc	o
719	atcgatcgagcgttctc	o
720	bgagaacgctcgaccttccat	sos
721	gctagacgcttagcgtga	sos
722	atcgactctcgagcgttctc	o
723	tagacgcttagcgtga	o
724	cgactctcgagcgttctc	sos
725	ggggtcgaccttggagggggg	o
726	gctaacgcttagcgtga	o
727	cgtcgtcgt	o
728	gagaacgctggacztccat	o
729	atcgacctacgtgcgttztc	o
730	atzgacctacgtgcgttctc	o
731	gctagazgcttagcgt	o
732	atcgactctcgagzggttctc	sos
733	ggggtaatgcatcagggggg	s
734	ggctgtattcctgactgcc	o
735	ccatgctaacctctagc	o
736	gctagatgcttagcgtga	o
737	cgtaccttacggtga	o
738	tccatgctggctcctgatgct	o
739	atcgactctctcgagcgttctc	o
740	gctagagcttagcgtga	o
741	atcgactctcgagtggttctc	o
742	aacgctcgaccttccat	o
743	ctcaacgctggaccttccat	o
744	atcgacctacgtgcgttctc	o
745	gagaatgctggaccttccat	o
746	tcacgctaacctctgac	o

10017695.121801

747	bgagaacgctccagcactgat	o
748	bgagcaagctggaccttccat	o
749	cgctagagggttagcgtga	o
750	gctagatgttaacgt	o
751	atggaaggtccacgttctc	o
752	gctagatgttagcgt	o
753	gctagacgttagtgt	o
754	tccatgacggctcctgatgct	o
755	tccatggcggctcctgatgct	o
756	gctagacgatagcgt	o
757	gctagtgcgatagcgt	o
758	tccatgacgttctcctgatgct	o
759	tccatgtcgttctcctgatgct	o
760	gctagacgttagzgt	o
761	gctaggcgttagcgt	o
762	tccatgtzggctcctgatgct	o
763	tccatgtcggctzctgatgct	o
764	atzgactctzgagzgttctc	o
765	atggaaggtccagtgttctc	o
766	gcatgacgttgagct	s
767	ggggtcaacgttgagggggg	sos
768	ggggtcaagtctgagggggg	o
769	cgcgcgcgcgcgcgcgcgcg	s
770	cccccccccccccccccccccccc	s
771	cccccccccccccccccccccccc	o
772	tccatgtcgtcctgatcct	o
773	gctaaacgttagcgt	o
774	tccatgtcgatcctgatgct	o
775	tccatgccggctcctgatgct	sos
776	aaaatcaacgttgaaaaaaa	o
777	tccataacgttcctgatgct	o
778	tggaggtcccaccgagatcgag	s
779	cgtcgtcgtcgtcgtcgtcgt	s
780	ctgctgctgctgctgctgctg	s
781	gagaacgctccgaccttcgat	s
782	gctagatgttagcgt	s
783	gcatgacgttgagct	o
784	tcaatgctgaf	o
785	tcaacgttgaf	o
786	tcaacgttgab	o
787	gcaatattgcb	o
788	gcaatattgcf	o
789	agttgcaact	o
790	tcttcgaa	o
791	tcaacgtc	o
792	ccatgtcggctcctgatgct	o
793	gtttttatataatttggg	o
794	tttttgtttgctgcttttgctgtt	s
795	ttgggggggggtt	s
796	ggggttgggggtt	o
797	ggtggtgtaggttttgg	o
798	bgagaazgctcgaccttcgat	o
799	tcaacgttaacgttaacgtt	o
800	bgagcaagztggaccttccat	o
801	bgagaazgctccagcactgat	o
802	tcaazgttgax	o
803	gzaatattgcx	o

747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803

804	tgctgcttttgcgcttttgcgctt	o
805	ctgcgcttagcaatttaactgtg	o
806	tccatgacgttcctgatgct	s
807	tgcattgccgtgcatccgtacacagctct	s
808	tgcattgccgtacacagctct	s
809	tgcattcagctct	s
810	tgcgctct	s
811	cccccccccccccccccccc	s
812	cccccccccccc	s
813	cccccccc	sos
814	tgcattcagctct	o
815	tgcattgccgtacacagctct	s
816	gagcaagctggaccttccat	s
817	tcaacgttaacgttaacgttaacgttaacgtt	s
818	gagaacgctcgaccttcgat	o
819	gtccccatttcccagaggaggaaat	o
820	ctagcggctgacgtcatcaagctag	o
821	ctagcttgatgacgtcagccgctag	s
822	cggctgacgtcatcaa	o
823	ctgacgtg	o
824	ctgacgtcat	o
825	attcgatcggggcgggcgag	o
826	ctcgccccgccccgatcgaat	o
827	gactgacgtcagcgt	s
828	ctagcggctgacgtcataaagctagc	s
829	ctagctttatgacgtcagccgctagc	s
830	ctagcggctgagctcataaagctagc	s
831	ctagtggctgacgtcatcaagctag	s
832	tccaccacgtggctcatgct	o
833	gggaatgaaagattttattataag	o
834	tctaaaaaccatctattcttaaccct	o
835	agctcaacgtcatgc	o
836	ttaacggtggttagcgggtattggctc	o
837	ttaagaccaataaccgctaccaccg	o
838	gatctagtgatgagtcagccggatc	o
839	gatccggctgactcatcactagatc	o
840	tccaagacgttcctgatgct	o
841	tccatgacgtccctgatgct	o
842	tccaccacgtggctgatgct	o
843	ccacgtggacctctagc	o
844	tcagaccacgtggctcgggtgttctga	o
845	tcaggaacacccgaccacgtggctctga	o
846	catttccacgatttccca	o
847	ttctctctgcaagagact	o
848	tgtatctctctgaaggact	o
849	ataaagcgaaactagcagcagtttc	o
850	gaaactgctgctagtttgcgtttat	o
851	tgcccaaagaggaaaatttggttcatacag	o
852	ctgtatgaaacaaatttctctttgggca	s
853	ttaggggttaggggttaggggt	s
854	tccatgagcttcctgatgct	s
855	aaaacatgacgttcaaaaaa	s
856	aaaacatgacgttcggggggg	sos
857	ggggcatgagcttcggggggg	o
858	ctaggctgacgtcatcaagctag	o
859	tctgacgtcatctgacgttggtgacgtct	o
860	ggaattagtaatatagataagaagtt	o

100155-1001

861	tttacctttttataaacataactaaaacaaa	o
862	gcgttttttttttgcg	s
863	atatctaatacaaaacattaacaaa	o
864	tctatcccagggtggttcctgttag	o
865	btccatgacgttcctgatgct	o
866	btccatgacgttcctgatgct	o
867	ttttttttttttttf	so
868	ttttttttttttttf	o
869	ctagcttgatgagctcagccgctag	o
870	ttcagttgtcttgcgttagctaa	s
871	tccatgagcttcctgagtct	o
872	ctagcggctgacgtcatcaatctag	s
873	tgctagctgtgcctgtacct	o
874	atgctaaaggacgtcacattgca	o
875	tgcaatgtgacgtccttttagcat	o
876	gtaggggactttccgagctcgagatcctatg	o
877	cataggatctcgagctcggaaagtcccctac	o
878	ctgtcaggaactgcaggttaagg	o
879	cataacataggaatatctactcctcgc	o
880	ctccagctccaagaaaggacg	o
881	gaagtttctggtaagtcttcg	s
882	tgctgcttttgcgttttgcgtt	s
883	tcgtcgtttgcgttttgcgtt	s
884	tcgtcgtttgcgttttgcgtt	s
885	tcctgacgttcggcgcgcgccc	
886	tgctgcttttgcgttttgcgtt	s
887	tccatgagcttcctgagctt	s
888	tcgtcgtttgcgttcgttttgcgtt	s
889	tcgtcgtttgcgttcgttttgcgtt	s
890	tcgcgtgcgttttgcgttttgcgtt	s
891	ttcgtcgttttgcgttttgcgtt	s
892	tcctgacggggaagt	s
893	tcctggcgtggaagt	s
894	tcctggcgtggaagt	s
895	tcctggcgtggaagt	s
896	tcctgacgtggaagt	s
897	gcgacgttcggcgcgcgccc	s
898	gcgacgggcgcgcgcgccc	s
899	gcggcgtgcggcgcgcgccc	s
900	gcggcgtgcggcgcgcgccc	s
901	gcgacgttcggcgcgcgccc	s
902	gcggcgttcggcgcgcgccc	s
903	gcgacgtgcggcgcgcgccc	s
904	tcgtcgctgtctccg	s
905	tgtgggggttttggttttg	s
906	aggggaggggaggggagggg	s
907	tgtgtgtgtgtgtgtgtgt	sos
908	ctctctctctctctctctct	s
909	ggggtcgacgtcgagggggg	s
910	atatatatatatatatatat	s
911	tttttttttttttttttttttt	s
912	tttttttttttttttttttt	s
913	tttttttttttttttttt	
914	gctagaggggaggggt	
915	gctagatgttagggg	
916	gcatgagggggagct	
917	atggaaggtccaggggggctc	

100795-1301

100795100

918	atggactctggagggggctc	
919	atggaaggtccaaggggctc	
920	gagaaggggggaccttggat	
921	gagaaggggggaccttccat	
922	gagaagggggccagcactgat	
923	tccatgtggggcctgatgct	
924	tccatgaggggcctgatgct	
925	tccatgtggggcctgatgct	
926	atggactctccggggttctc	
927	atggaaggtccggggttctc	
928	atggactctggaggggtctc	
929	atggaggtccatggggctc	
930	atggactctggggggttctc	
931	tccatgtgggtggggatgct	
932	tccatgcgggtggggatgct	
933	tccatgggggtcctgatgct	
934	tccatgggggtcctgatgct	
935	tccatgggggtgctgatgct	
936	tccatgggggttctgatgct	
937	tccatcgggggcctgatgct	
938	gctagaggagtgct	s
939	tttttttttttttttttt	s
940	gmgtcaacgttgagggmggg	s
941	ggggagttcggtgaggggggg	s
942	tcgtcggtttcccccccccc	s
943	ttggggggttttttttttttt	s
944	tttaaattttaaaatttaaata	s
945	ttggtttttttggttttttttg	s
946	tttcccttttcccttttccctc	sos
947	ggggtcatcgatgagggggg s	
948	tccatgacgttcctgacgtt	
949	tccatgacgttcctgacgtt	
950	tccatgacgttcctgacgtt	
951	tccatgacgttcctgacgtt	
952	tccatgacgttcctgacgtt	
953	tccatgacgttcctgacgtt	
954	tccatgacgttcctgacgtt	
955	tccatgacgttcctgacgtt	
956	tccatgacgttcctgacgtt	
957	tccatgacgttcctgacgtt	
958	tccatgacgttcctgacgtt	sos
959	gggggacgatcgctcggggg	sos
960	gggggtcgtagcagggggg	po
961	tttttttttttttttttttt	po
962	aaaaaaaaaaaaaaaaaaaaaa	po
963	cccccccccccccccccccc	
964	tcgtcggtttgtcggtttgtcggt	
965	tcgtcggtttgtcggtttgtcggt	
966	tcgtcggtttgtcggtttgtcggt	
967	tcgtcggtttgtcggtttgtcggt	
968	gggggtcaacgttgagggggg	
969	gggggtcaacgttgagggggg	
970	gggggtcaagcttgagggggg	
971	tgtgttcccccccccccc	

10017995-1001

972	ggggacgtcgacgtggggggg	sos
973	gggggtcgtcgacgagggggg	sos
974	gggggtcgacgtacgtcgagggggg	sos
975	ggggaccggtaccggtggggggg	sos
976	gggtcgacgtcgaggggggg	sos
977	gggggtcgacgtcgagggggg	sos
978	ggggaacgttaacgttggggggg	sos
979	gggggtcacggtgaggggggg	sos
980	gggggtcgttcgaacgagggggg	sos
981	ggggacgttcgaacgtggggggg	s
982	tcaactttga	s
983	tcaagcttga	s
984	tcacgatcgtga	s
985	tcagcatgctga	sos
986	ggggggagcatgctgggggggg	sos
987	ggggggggggggggggggggg	sos
988	ggggggacgatatcgctcgggggg	sos
989	ggggggacgacgtcgctcgggggg	sos
990	ggggggacgagctcgctcgggggg	sos
991	ggggggacgtacgtcgggggg	
992	tcaacgtt	
993	tccataccggtcctgatgct	s
994	tccataccggtcctaccggt	sos
995	ggggggacgatcgttggggggg	sos
996	ggggaacgatcgctcgggggg	sos
997	ggg ggg acg atc gtc ggg ggg	sos
998	ggg gga cga tcg tcg ggg ggg	po
999	aaa gac gtt aaa	po
1000	aaagagcttaaa	po
1001	aaagazgttaaa	po
1002	aaattcggaaaa	sos
1003	gggggtcatcgatgagggggg	sos
1004	gggggtcaacgttgagggggg	po
1005	atgtagcttaataacaaagc	po
1006	ggatcccttgagttacttct	po
1007	ccattccacttctgattacc	po
1008	tatgtattatcatgtagata	po
1009	agcctacgtattcacccctcc	po
1010	ttcctgcaactactattgta	po
1011	atagaaggccctacaccagt	po
1012	ttacaccggtctatggaggt	po
1013	ctaaccagatcaagtctagg	po
1014	cctagacttgatctggttag	po
1015	tataagcctcgccgacatg	po
1016	catgtcggacgaggttata	po
1017	tggtggtggggagtaagctc	po
1018	gagctactccccaccacca	po
1019	gccttcgatcttcgttggga	po
1020	tggacttctctttgccgtct	po
1021	atgctgtagcccagcgataa	

1022	accgaatcagcggaaagtga	po
1023	tccatgacgttcctgacgtt	
1024	ggagaaacccatgagctcatctgg	
1025	accacagaccagcaggcaga	
1026	gagcgtgaactgcgcgaaga	
1027	tcggtacccttgacgcggtt	
1028	ctggagccctagccaaggat	
1029	gcgactccatcaccagcgat	
1030	cctgaagtaagaaccagatgt	
1031	ctgtgttatctgacatacacc	
1032	aattagccttaggtgattggg	
1033	acatctggttcttacttcagg	
1034	ataagtcataatcttgggaactac	
1035	ccaatcacctaaggctaatt	sos
1036	ggggtcgtcgacgagggggg	sos
1037	ggggtcgttcgaacgagggggg	sos
1038	ggggacgttcgaacgtgggggg	s
1039	tcctggcgqggaagt	sos
1040	ggggaacgacgtcggtgggggg	sos
1041	ggggaacgtacgtcgggggg	sos
1042	ggggaacgtacgtacgttgggggg	sos
1043	ggggtcacgggtgagggggg	sos
1044	ggggtcgacgtacgtcgagggggg	sos
1045	ggggaccgggtaccgggtgggggg	sos
1046	gggtcgacgtcgagggggg	sos
1047	ggggtcgacgtcgagggg	sos
1048	ggggaacgttaacgttgggggg	sos
1049	ggggacgtcgacgtggggg	
1050	gcactcttcgaagctacagccggcagcctctgat	
1051	cggctcttccatgaggtctttgctaactcttg	
1052	cggctcttccatgaaagtcttggacgatgtgagc	s
1053	tcctgcaggttaagt	sos
1054	gggggtcgttcgttgggggg	sos
1055	gggggatgattggttgggggg	sos
1056	gggggazgatzggttgggggg	sos
1057	gggggagctagcttgggggg	s
1058	ggttcttttggtccttgtct	s
1059	ggttcttttggtcctcgctct	s
1060	ggttcttttggtccttatct	s
1061	ggttcttgggttcttctgtct	s
1062	tggtcttttggtccttgtct	s
1063	ggttcaaatggtccttgtct	s
1064	gggtcttttgggccttgtct	s
1065	tccaggacttctctcaggtttttt	s
1066	tccaaaacttctctcaaat	s
1067	tactacttttatacttttatactt	s
1068	tgtgtgtgtgtgtgtgtgtgtg	s
1069	ttgttgttgttgttgttgttgtg	s
1070	ggctccggggaggggaatcttctctat	sos
1071	gggacgatcggtcggggggg	

1007995-1331

1072	gggtcgtcgacgaggggggg	sos
1073	ggtcgtcgacgaggggggg	sos
1074	gggtcgtcgtcgtggggggg	sos
1075	ggggacgatcgtcggggggg	sos
1076	ggggacgtcgtcgtgggggg	sos
1077	ggggtcgacgtcgacgtcgagggggg	sos
1078	ggggaaccgcggttggggggg	sos
1079	ggggacgacgtcgtggggggg	sos
1080	tcgtcgtcgtcgtcgtggggggg	s
1081	tcctgccggggaagt	s
1082	tcctgcaggggaagt	s
1083	tcctgaaggggaagt	s
1084	tcctggcgggcaagt	s
1085	tcctggcgggtaagt	s
1086	tcctggcgggaaagt	s
1087	tcggggcggggaagt	s
1088	tcggggcggggaagt	s
1089	tccgggcggggaagt	s
1090	gggggacgttggggg	sos
1091	gggggttttttttttgggggg	sos
1092	ggggccccccccccgggggg	sos
1093	gggggttggtggttgggggg	sos

Nucleic acids having modified backbones also are included in the class of nucleic acids having antiangiogenic properties. Modified backbone nucleic acids include those having phosphorothioate, methylphosphonate, methylphosphorothioate, p-ethoxy and/or phosphorodithioate internucleotide or internucleoside bonds. Chimeric oligonucleotides having mixtures of modified and/or unmodified backbones also are included in the invention.

In the case when an antiangiogenic nucleic acid is administered in conjunction with a nucleic acid vector, it is preferred that the backbone of the antiangiogenic nucleic acid be a chimeric combination of phosphodiester and phosphorothioate bonds (or other modification of the internucleotide bonds). This is because the uptake of the plasmid vector by the cell may be hindered by the presence of completely phosphorothioate oligonucleotide. Thus when both a vector and an oligonucleotide are delivered to a subject, it is preferred that the oligonucleotide have chimeric or phosphorothioate internucleotide bonds and that the plasmid be associated with a vehicle that delivers it directly into the cell, thus avoiding the need for cellular uptake. Such vehicles are known in the art and include, for example, liposomes, electroporation devices and gene guns.

For use in the instant invention, the antiangiogenic nucleic acids can be synthesized *de novo* using any of a number of procedures well known in the art. Such compounds are

referred to as "synthetic nucleic acids." For example, the b-cyanoethyl phosphoramidite method (Beaucage, S.L., and Caruthers, M.H., *Tet. Let.* 22:1859, 1981); nucleoside H-phosphonate method (Garegg *et al.*, *Tet. Let.* 27:4051-4054, 1986; Froehler *et al.*, *Nucl. Acid. Res.* 14:5399-5407, 1986; Garegg *et al.*, *Tet. Let.* 27:4055-4058, 1986; Gaffney *et al.*, *Tet. Let.* 29:2619-2622, 1988). These chemistries can be performed by a variety of automated oligonucleotide synthesizers available in the market.

Alternatively, nucleic acids can be produced on a large scale in plasmids, (see, e.g., Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York, 1989) and separated into smaller pieces or administered whole. Nucleic acids can be prepared from existing nucleic acid sequences (e.g., genomic or cDNA) using known techniques, such as those employing restriction enzymes, exonucleases or endonucleases. Nucleic acids prepared in this manner are referred to as isolated nucleic acids. The term "antiangiogenic nucleic acid" encompasses both synthetic and isolated antiangiogenic nucleic acids.

For use *in vivo*, nucleic acids are preferably relatively resistant to degradation (e.g., are stabilized). A "stabilized nucleic acid molecule" as used herein means a nucleic acid molecule that is relatively resistant to *in vivo* degradation (e.g. via an exo- or endo-nuclease). Stabilization can be a function of length or secondary structure. Antiangiogenic nucleic acids that are tens to hundreds of kilobases long are relatively resistant to *in vivo* degradation. For shorter antiangiogenic nucleic acids, secondary structure can stabilize and increase their effect. For example, if the 3' end of a nucleic acid is self-complementary to an upstream region of the same nucleic acid, so that it can fold back and form a stem/loop structure by internal self-hybridization, then the nucleic acid may be stabilized and therefore may exhibit more *in vivo* activity.

Alternatively, nucleic acid stabilization can be accomplished via backbone modifications. Preferred stabilized nucleic acids of the instant invention have a modified backbone. It has been demonstrated that modification of the nucleic acid backbone provides enhanced activity of the antiangiogenic nucleic acids when administered *in vivo*. One type of modified backbone is a phosphate backbone modification. For example, antiangiogenic nucleic acids including at least two phosphorothioate linkages at the 5' end of the oligonucleotide and multiple phosphorothioate linkages at the 3' end, preferably 5 or more, can in some circumstances protect the nucleic acid from degradation by intracellular exo- and endo-nucleases and thereby provide maximal activity. Other phosphate modified nucleic

acids include phosphodiester modified nucleic acids, combinations of phosphodiester and phosphorothioate nucleic acids, methylphosphonate, methylphosphorothioate, phosphorodithioate, p-ethoxy and combinations thereof. Some of these combinations in CpG nucleic acids and their particular effects on immune cells is discussed in more detail in PCT Published Patent Applications PCT/US95/01570 and PCT/US97/19791, the entire contents of which are hereby incorporated by reference. Although not intending to be bound by any particular theory, it is believed that these modified nucleic acids may have increased activity relative to unmodified nucleic acids due to enhanced nuclease resistance, increased cellular uptake, increased protein binding, and/or altered intracellular localization.

Modified backbone nucleic acids, such as those having phosphorothioates bonds may be synthesized using automated techniques employing, for example, phosphoramidate or H-phosphonate chemistries. Aryl- and alkyl-phosphonates can be made, e.g., as described in U.S. Patent No. 4,469,863. Alkylphosphotriesters, in which the charged oxygen moiety is alkylated as described in U.S. Patent No. 5,023,243 and European Patent No. 092,574, can be prepared by automated solid phase synthesis using commercially available reagents. Methods for making other nucleic acid backbone modifications and substitutions have been described (Uhlmann, E. and Peyman, A., *Chem. Rev.* 90:544, 1990; Goodchild, J., *Bioconjugate Chem.* 1:165, 1990).

Another type of modified backbone, useful according to the invention, is a peptide nucleic acid. The backbone is composed of aminoethylglycine and supports bases which provide the nucleic acid character. The backbone does not include any phosphate and thus may optionally have no net charge. The lack of charge allows for stronger DNA-DNA binding because the charge repulsion between the two strands does not exist. Additionally, because the backbone has an extra methylene group, the oligonucleotides are enzyme/protease resistant. Peptide nucleic acids can be purchased from various commercial sources, e.g., Perkin Elmer, or synthesized de novo.

Another class of backbone modifications include 2'-O-methylribonucleosides (2'-O-Me). These types of substitutions are described extensively in the literature and in particular with respect to their immunostimulating properties in Zhao et al., *Bioorganic and Medicinal Chemistry Letters*, 1999, 9:24:3453. Zhao et al. describes methods of preparing 2'-O-Me modifications to nucleic acids.

The nucleic acid molecules of the invention may include naturally-occurring or synthetic purine or pyrimidine heterocyclic bases as well as modified backbones. Purine or

pyrimidine heterocyclic bases include, but are not limited to, adenine, guanine, cytosine, thymidine, uracil, and inosine. Other representative heterocyclic bases are disclosed in US Patent No. 3,687,808, issued to Merigan, et al. The terms "purines" or "pyrimidines" or "bases" are used herein to refer to both naturally-occurring or synthetic purines, pyrimidines or bases.

Other stabilized nucleic acids include non-ionic DNA analogs, such as alkyl- and aryl-phosphates (in which the charged phosphonate oxygen is replaced by an alkyl or aryl group), phosphodiester and alkylphosphotriesters, in which the charged oxygen moiety is alkylated. Nucleic acids which contain diol, such as tetraethyleneglycol or hexaethyleneglycol, at either or both termini have also been shown to be substantially resistant to nuclease degradation.

The antiangiogenic nucleic acids having backbone modifications useful according to the invention in some embodiments are S- or R-chiral antiangiogenic nucleic acids. An "S chiral antiangiogenic nucleic acid" as used herein is an antiangiogenic nucleic acid wherein at least two nucleotides have a backbone modification forming a chiral center and wherein a plurality of the chiral centers have S chirality. An "R chiral antiangiogenic nucleic acid" as used herein is an antiangiogenic nucleic acid wherein at least two nucleotides have a backbone modification forming a chiral center and wherein a plurality of the chiral centers have R chirality. The backbone modification may be any type of modification that forms a chiral center. The modifications include but are not limited to phosphorothioate, methylphosphonate, methylphosphorothioate, phosphorodithioate, p-ethoxy, 2'-O-Me and combinations thereof.

The chiral antiangiogenic nucleic acids must have at least two nucleotides within the nucleic acid that have a backbone modification. All or less than all of the nucleotides in the nucleic acid, however, may have a modified backbone. Of the nucleotides having a modified backbone (referred to as chiral centers), a plurality have a single chirality, S or R. A "plurality" as used herein refers to an amount greater than 50%. Thus, less than all of the chiral centers may have S or R chirality as long as a plurality of the chiral centers have S or R chirality. In some embodiments at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of the chiral centers have S or R chirality. In other embodiments at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of the nucleotides have backbone modifications.

The S- and R- chiral antiangiogenic nucleic acids may be prepared by any method known in the art for producing chirally pure oligonucleotides. Stec et al teach methods for

producing stereopure phosphorothioate oligodeoxynucleotides using an oxathiaphospholane. (Stec, W.J., et al., 1995, *J. Am. Chem. Soc.*, 117:12019). Other methods for making chirally pure oligonucleotides have been described by companies such as ISIS Pharmaceuticals. US Patents which disclose methods for generating stereopure oligonucleotides include 5,883,237; 5,837,856; 5,599,797; 5,512,668; 5,856,465; 5,359,052; 5,506,212; 5,521,302; and 5,212,295, each of which is hereby incorporated by reference in its entirety.

As used herein, administration of an antiangiogenic nucleic acid is intended to embrace the administration of one or more antiangiogenic nucleic acids which may or may not differ in terms of their profile, sequence, backbone modifications and biological effect. As an example, CpG nucleic acids and T-rich nucleic acids may be administered to a single subject along with other antiangiogenic medicament(s), such as endostatin or angiostatin. In another example, a plurality of CpG nucleic acids which differ in nucleotide sequence may also be administered to a subject.

The invention encompasses the administration of the antiangiogenic nucleic acids along with other medicaments in order to provide a synergistic effect useful in the prevention and/or treatment of conditions that involve unwanted angiogenesis, such as cancer. Accordingly, methods for inhibition of angiogenesis are provided. The methods include the administration of at least one antiangiogenic nucleic acid formulated for administration to a subject. Non-nucleic acid antiangiogenesis molecules also can be administered to the subject, including, but not limited to endogenous angiogenesis inhibitors including PD 174073 and PD 166285 (Parke-Davis), SU5416 and SU6668 (Sugen), ZD 4190 and ZD 6474 (Zeneca), PTK 787 (also known as CGP79787 or ZK22584) (Novartis), Anti-VEGF mAb (Genentech), Anti-KDR mAb (ImClone), RPI 4610 (Ribozyme), TNP 470 (Abbott/TAP), AG 3340 (Agouron), Marimastat (British Biotech), Bay 12-9566 (Bayer), Neovastat (Aeterna), BMS 275291 (Bristol Myers-Squibb), CGS 27023A (Novartis), D1927 Chiroscience), D2163 (Chiroscience), Isoquinolines (Pfizer), Vitaxin (IXSYS), S-137 (Searle), S-836 (Searle), SM256 (Dupont), SG545 (Dupont), Angiostatin (EntreMed), Endostatin (EntreMed), Thalidomide (EntreMed), Squalamine (Magainin), CAI (National Cancer Institute), CM-101 (CarboMed), U-995 (Gwo-Chyang GMP), Combretastatin A-4 (Oxigene), platelet factor-4, vasostatin, thrombospondin, tissue inhibitors of metalloproteinases (TIMPs), STI412 (Sun and McMahon, *Drug Discov. Today* 5(8):344-353, 2000; Klohs and Hamby, *Curr. Opin. Biotechnol.* 10:544-549, 1999), fumagillin, non-glucocorticoid steroids and heparin and heparin fragments and antibodies to one or more angiogenic peptides such as α -FGF, β -FGF,

VEGF, IL-8, and GM-CSF. Some of the foregoing may be administered in the form of nucleic acids encoding proteins; in each case the active agent is a protein and not the nucleic acid encoding the protein.

The antiangiogenic nucleic acid molecules of the invention can be administered concurrently with, or sequentially with, the non-nucleic acid antiangiogenesis molecules described above. Coadministration may be in the form of administration of a composition containing both kinds of antiangiogenic agents, or a plurality of compositions, each of which may contain one or more than one of the antiangiogenic agents.

The invention may be used in the treatment of cancer, but is not so limited. In these methods, an effective amount of at least one antiangiogenic nucleic acid is administered to a subject having cancer, or in other instances a subject at risk of developing cancer. Other non-nucleic acid antiangiogenesis molecules also can be administered, as described above. In addition, in certain embodiments of the invention, anticancer molecules are administered in combination with the antiangiogenesis molecules.

The compounds useful in the invention may be delivered in a mixture with anti-proliferative agents (particularly anticancer agents) which are not antiangiogenic nucleic acids. One of ordinary skill in the art is familiar with a variety of anti-proliferative agents which are used in the medical arts to treat proliferative diseases such as cancer. These anti-cancer agents may act by directly killing cells, such as cancer cells (i.e., direct action anti-cancer agents), or alternatively they may act by sensitizing cells to direct action anti-cancer agents (i.e., indirect action anti-cancer agents). Those of skill in the art will recognize the distinction and are familiar with agents of either class. Anticancer agents include, but are not limited to, the following sub-classes of compounds:

Antineoplastic agents such as: Acivicin; Aclarubicin; Acodazole Hydrochloride; Acronine; Adozelesin; Adriamycin; Aldesleukin ; Altretamine; Ambomycin; Ametantrone Acetate; Aminoglutethimide; Amsacrine; Anastrozole; Anthramycin; Asparaginase; Asperlin ; Azacitidine; Azetepa; Azotomycin; Batimastat; Benzodepa; Bicalutamide; Bisantrone Hydrochloride; Bisnafide Dimesylate; Bizelesin; Bleomycin Sulfate; Brequinar Sodium; Bropiramine ; Busulfan; Cactinomycin; Calusterone; Caracemide; Carbetimer; Carboplatin; Carmustine; Carubicin Hydrochloride; Carzelesin; Cedefingol; Chlorambucil; Cirolemycin ; Cisplatin; Cladribine; Crisnatol Mesylate; Cyclophosphamide ; Cytarabine ; Dacarbazine; DACA (N-[2-(Dimethyl-amino)ethyl]acridine-4-carboxamide); Dactinomycin; Daunorubicin Hydrochloride; Daunomycin; Decitabine; Dexormaplatin; Dezaguanine; Dezaguanine

Mesylate; Diaziquone; Docetaxel; Doxorubicin; Doxorubicin Hydrochloride; Droloxifene;
 Droloxifene Citrate; Dromostanolone Propionate; Duazomycin; Edatrexate; Eflornithine
 Hydrochloride ; Elsamitucin; Enloplatin; Enpromate; Epiropidine; Epirubicin
 Hydrochloride; Erbulozole; Esorubicin Hydrochloride; Estramustine; Estramustine Phosphate
 5 Sodium; Etanidazole; Ethiodized Oil I 131; Etoposide; Etoposide Phosphate; Etoprine;
 Fadzole Hydrochloride; Fazarabine; Fenretinide; Floxuridine; Fludarabine Phosphate;
 Fluorouracil; 5-FdUMP; Flucitabine; Fosquidone; Fostriecin Sodium; Gemcitabine;
 Gemcitabine Hydrochloride; Gold Au 198 ; Hydroxyurea; Idarubicin Hydrochloride;
 Ifosfamide; Ilmofofosine; Interferon Alfa-2a ; Interferon Alfa-2b ; Interferon Alfa-n1;
 Interferon Alfa-n3; Interferon Beta- I a ; Interferon Gamma- I b; Iproplatin; Irinotecan
 Hydrochloride ; Lanreotide Acetate; Letrozole; Leuprolide Acetate ; Liarozole Hydrochloride;
 Lometrexol Sodium; Lomustine; Losoxantrone Hydrochloride; Masoprocol; Maytansine;
 Mechlorethamine Hydrochloride; Megestrol Acetate; Melengestrol Acetate; Melphalan;
 Menogaril; Mercaptopurine; Methotrexate; Methotrexate Sodium; Metoprine; Meturedepa;
 10 Mitindomide; Mitocarcin; Mitocromin; Mitogillin; Mitomalcin; Mitomycin; Mitosper;
 Mitotane; Mitoxantrone Hydrochloride; Mycophenolic Acid; Nocodazole; Nogalamycin;
 Ormaplatin; Oxisuran; Paclitaxel; Pegaspargase; Peliomycin; Pentamustine; Peplomycin
 Sulfate; Perfosfamide; Pipobroman; Pipo sulfan; Piroxantrone Hydrochloride; Plicamycin;
 Plomestane; Porfimer Sodium; Porfiromycin ; Prednimustine; Procarbazine Hydrochloride;
 20 Puromycin; Puromycin Hydrochloride; Pyrazofurin; Riboprine; Rogletimide; Safingol ;
 Safingol Hydrochloride ; Semustine; Simtrazene; Sparfosate Sodium; Sparsomycin;
 Spirogermanium Hydrochloride; Spiromustine; Spiroplatin; Streptonigrin; Streptozocin;
 Strontium Chloride Sr 89; Sulofenur; Talisomycin; Taxane; Taxoid; Tecogalan Sodium;
 Tegafur; Teloxantrone Hydrochloride; Temoporfin; Teniposide; Teroxirone; Testolactone;
 25 Thiamiprine; Thioguanine; Thiotepa; Thymitaq; Tiazofurin; Tirapazamine; Tomudex; TOP-
 53; Topotecan Hydrochloride; Toremifene Citrate; Trestolone Acetate; Triciribine Phosphate;
 Trimetrexate; Trimetrexate Glucuronate; Triptorelin; Tubulozole Hydrochloride; Uracil
 Mustard; Uredepa; Vapreotide; Verteporfin; Vinblastine; Vinblastine Sulfate; Vincristine;
 Vincristine Sulfate; Vindesine; Vindesine Sulfate; Vinepidine Sulfate; Vinglycinat Sulfate;
 30 Vinleurosine Sulfate; Vinorelbine Tartrate; Vinrosidine Sulfate; Vinzolidine Sulfate;
 Vorozole; Zeniplatin; Zinostatin; Zorubicin Hydrochloride; 2-Chlorodeoxyadenosine; 2'-
 Deoxyformycin; 9-aminocamptothecin; raltitrexed; N-propargyl-5,8-dideazafolic acid; 2-

chloro-2'-arabino-fluoro-2'-deoxyadenosine; 2-chloro-2'-deoxyadenosine; anisomycin; trichostatin A; hPRL-G129R; CEP-751; linomide.

Other anti-neoplastic compounds include: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives (e.g., 10-hydroxy- camptothecin); canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrodidemnin B; deslorelin; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; discodermolide; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epothilones including desoxyepothilones (A, R = H; B, R = Me); epithilones; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide; etoposide 4'-phosphate (etopofos); exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam;

heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene;
idramantone; ilmofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant
peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons;
interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; irinotecan; iroplact; irsogladine;
5 isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N
triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole;
leukemia inhibiting factor; leukocyte alpha interferon; leuprolide + estrogen + progesterone;
leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide;
lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol;
10 lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline;
lytic peptides; maitansine; mannostatin A; marimastat; masoprocil; maspin; matrilysin
inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin;
methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim;
mismatched double stranded RNA; mithracin; mitoguazone; mitolactol; mitomycin
15 analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene;
molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid
A + myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple
tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial
cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin;
20 nagrestip; naloxone + pentazocine; napavin; naphterpin; nartograstim; nedaplatin;
nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide
modulators; nitroxide antioxidant; nitrullin; O6-benzylguanine; octreotide; okicenone;
oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer;
ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel analogues; paclitaxel derivatives;
25 palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin;
pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole;
perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase
inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B;
plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine
30 complex; podophyllotoxin; porfimer sodium; porfiromycin; propyl bis-acridone;
prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C
inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors;
purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated

hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thalidomide; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene dichloride; topotecan; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; zinostatin stimalamer.

Anti-cancer Supplementary Potentiating Agents: Tricyclic anti-depressant drugs (e.g., imipramine, desipramine, amitriptyline, clomipramine, trimipramine, doxepin, nortriptyline, protriptyline, amoxapine and maprotiline); non-tricyclic anti-depressant drugs (e.g., sertraline, trazodone and citalopram); Ca^{++} antagonists (e.g., verapamil, nifedipine, nitrendipine and caroverine); Calmodulin inhibitors (e.g., prenylamine, trifluoroperazine and clomipramine); Amphotericin B; Triparanol analogues (e.g., tamoxifen); antiarrhythmic drugs (e.g., quinidine); antihypertensive drugs (e.g., reserpine); Thiol depleters (e.g., buthionine and sulfoximine) and Multiple Drug Resistance reducing agents such as Cremaphor EL. The compounds of the invention also can be administered with cytokines such as granulocyte colony stimulating factor.

Antiproliferative agent: Piritrexim Isethionate.

Radioactive agents: Fibrinogen I 125 ; Fludeoxyglucose F 18 ; Fluorodopa F 18 ;
Insulin I 125; Insulin I 131; Iobenguane I 123; Iodipamide Sodium I 131 ; Iodoantipyrine I
131 ; Iodocholesterol I 131 ; Iodohippurate Sodium I 123 ; Iodohippurate Sodium I 125 ;
Iodohippurate Sodium I 131 ; Iodopyracet I 125 ; Iodopyracet I 131 ; Iofetamine
5 Hydrochloride I 123 ; Iomethin I 125 ; Iomethin I 131 ; Iothalamate Sodium I 125 ;
Iothalamate Sodium I 131 ; Iotyrosine I 131; Liothyronine I 125; Liothyronine I 131;
Merisoprol Acetate Hg 197; Merisoprol Acetate Hg 203; Merisoprol Hg 197 ;
Selenomethionine Se 75 ; Technetium Tc 99m Antimony Trisulfide Colloid; Technetium Tc
99m Bicisate ; Technetium Tc 99m Disofenin ; Technetium Tc 99m Etidronate ; Technetium
10 Tc 99m Exametazime ; Technetium Tc 99m Furifosmin ; Technetium Tc 99m Glucoptate ;
Technetium Tc 99m Lidofenin ; Technetium Tc 99m Mebrofenin ; Technetium Tc 99m
Medronate ; Technetium Tc 99m Medronate Disodium; Technetium Tc 99m Mertiatide ;
Technetium Tc 99m Oxidronate ; Technetium Tc 99m Pentetate; Technetium Tc 99m
Pentetate Calcium Trisodium; Technetium Tc 99m Sestamibi ; Technetium Tc 99m
15 Siboroxime ; Technetium Tc 99m Succimer ; Technetium Tc 99m Sulfur Colloid ;
Technetium Tc 99m Teboroxime ; Technetium Tc 99m Tetrofosmin ; Technetium Tc 99m
Tiatide; Thyroxine I 125; Thyroxine I 131; Tolpovidone I 131 ; Triolein I 125; Triolein I 131.

The present invention further includes nucleic acid molecules formulated into a
pharmaceutical composition for the inhibition of angiogenesis. The pharmaceutical
20 compositions of the invention include those suitable for oral, rectal, nasal, topical (including
buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular,
intravenous, intratumoral and intradermal) administration.

The nucleic acids are delivered in effective amounts. In general, the term "effective
amount" of a nucleic acid refers to the amount necessary or sufficient to realize a desired
25 biologic effect. Specifically, the effective amount is that amount that reduces the rate or
inhibits altogether angiogenesis. For instance, when the subject bears a tumor having a blood
supply, an effective amount is that amount which decreases or eliminates all together the
blood supply to the tumor. Additionally, an effective amount may be that amount which
prevents an increase or causes a decrease in new blood vessels, e.g., those vessels supplying a
30 tumor. The effective amount may vary depending upon whether the antiangiogenic nucleic
acid is used alone or in combination with other therapeutics, or in single or multiple dosages.
In some instances, it is envisioned that the combination of antiangiogenic nucleic acids with
other therapeutic agents (which are themselves not antiangiogenic nucleic acids) can result in

5

15

25

30

buffering agents, preservatives, compatible carriers, adjuvants, and optionally other therapeutic ingredients.

For use in therapy, an effective amount of the nucleic acid can be administered to a subject by any mode that delivers the nucleic acid to a subject. "Administering" the pharmaceutical composition of the present invention may be accomplished by any means known to the skilled artisan. Some routes of administration include but are not limited to oral, intranasal, intratracheal, inhalation, ocular, vaginal, rectal, parenteral (e.g. intramuscular, intradermal, intravenous, intratumoral or subcutaneous injection) and direct injection.

For oral administration, the compounds (i.e., antiangiogenic nucleic acid molecules and optionally other antiangiogenesis agents) can be delivered alone without any pharmaceutical carriers or formulated readily by combining the active compound(s) with pharmaceutically acceptable carriers well known in the art. The term "pharmaceutically-acceptable carrier" means one or more compatible solid or liquid filler, diluents or encapsulating substances which are suitable for administration to a human or other vertebrate animal. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being commingled with the compounds of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficiency.

Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a subject to be treated. Pharmaceutical preparations for oral use can be obtained as solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Optionally the oral formulations may also be formulated in saline or buffers for neutralizing internal acid conditions.

Dragee cores may be provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl

pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

5 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or
10 suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. Microspheres formulated for oral administration may also be used. Such microspheres have been well defined in the art. All formulations for oral administration should be in dosages suitable for such administration.

15 For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

20 For administration by inhalation, the compounds for use according to the present invention may be conveniently delivered in the form of an aerosol spray, from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the
20 case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

25 The compounds, when it is desirable to deliver them systemically, may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

30 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as

ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active compounds may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal or vaginal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Suitable liquid or solid pharmaceutical preparation forms are, for example, aqueous or saline solutions for inhalation, microencapsulated, encochleated, coated onto microscopic gold particles, contained in liposomes, nebulized, aerosols, pellets for implantation into the skin, or dried onto a sharp object to be scratched into the skin. The pharmaceutical compositions may also include granules, powders, tablets, coated tablets, (micro)capsules, suppositories, syrups, emulsions, suspensions, creams, drops or preparations with protracted release of active compounds, in whose preparation excipients and additives and/or auxiliaries such as disintegrants, binders, coating agents, swelling agents, lubricants, flavorings, sweeteners or solubilizers are customarily used as described above. The pharmaceutical compositions are suitable for use in a variety of drug delivery systems. For a brief review of present methods for drug delivery, see Langer, *Science* 249:1527-1533, 1990, which is incorporated herein by reference.

The nucleic acid molecules and/or agents (*e.g.*, antiangiogenesis agents, anticancer agents) may be administered per se (neat) or in the form of a pharmaceutically acceptable salt.

When used in medicine the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically acceptable salts thereof. Such salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene sulphonic, tartaric, citric, methane sulphonic, formic, malonic, succinic, naphthalene-2-sulphonic, and benzene sulphonic. Also, such salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts of the carboxylic acid group.

Suitable buffering agents include: acetic acid and a salt (1-2% w/v); citric acid and a salt (1-3% w/v); boric acid and a salt (0.5-2.5% w/v); and phosphoric acid and a salt (0.8-2% w/v). Suitable preservatives include benzalkonium chloride (0.003-0.03% w/v); chlorobutanol (0.3-0.9% w/v); parabens (0.01-0.25% w/v) and thimerosal (0.004-0.02% w/v).

The nucleic acids or other therapeutics useful in the invention may be delivered in mixtures with additional antiangiogenesis agent(s). A mixture may consist of several antiangiogenesis agents in addition to the nucleic acid.

A variety of administration routes are available. The particular mode selected will depend, of course, upon the particular nucleic acid molecules or other agents selected, the particular condition being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of an immune response without causing clinically unacceptable adverse effects. Preferred modes of administration are discussed above.

The compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the compounds into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the compounds into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product. Liquid dose units are vials or ampoules. Solid dose units are tablets, capsules and suppositories.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the compounds, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer

base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di-and tri-glycerides; hydrogel release systems; systatic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which an agent of the invention is contained in a form within a matrix such as those described in U.S. Patent Nos. 4,452,775, 4,675,189, and 5,736,152, and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation. In still other embodiments, the agents and nucleic acids are formulated with GELFOAM, a commercial product consisting of modified collagen fibers that degrade slowly.

The nucleic acid may be directly administered to the subject or may be administered in conjunction with a pharmaceutically acceptable carrier or a delivery vehicle. The nucleic acid and optionally other therapeutic agents may be administered alone (e.g. in saline or buffer) or using any delivery vehicles known in the art. One type of delivery vehicle is referred to herein as a nucleic acid delivery complex. A "nucleic acid delivery complex" shall mean a nucleic acid molecule associated with (e.g. ionically or covalently bound to; or encapsulated within) a targeting means (e.g. a molecule that results in higher affinity binding to target cell (e.g. dendritic cell surfaces and/or increased cellular uptake by target cells). Examples of nucleic acid delivery complexes include nucleic acids associated with: a sterol (e.g. cholesterol), a lipid (e.g. a cationic lipid, virosome or liposome), or a target cell specific binding agent (e.g. a ligand recognized by target cell specific receptor). Preferred complexes may be sufficiently stable *in vivo* to reduce significant uncoupling prior to internalization by the target cell. However, the complex may be cleavable under appropriate conditions within the cell so that the nucleic acid may be released in a functional form.

The nucleic acid molecules may be delivered by non-invasive methods as described above. Non-invasive delivery of compounds is desirable for treatment of children, elderly, animals, and even adults and also to avoid the risk of needle-stick injury. Delivery vehicles for delivering compounds to mucosal surfaces have been described and include but are not

limited to: cochleates, emulsomes, ISCOMs, liposomes, live bacterial vectors (e.g., *Salmonella*, *Escherichia coli*, *Bacillus calmette-guerin*, *Shigella*, *Lactobacillus*), live viral vectors (e.g., Vaccinia, adenovirus, Herpes Simplex), microspheres, nucleic acid vaccines, polymers (e.g. carboxymethylcellulose, chitosan), polymer rings, proteosomes, sodium fluoride, transgenic plants, virosomes, and virus-like particles.

Examples

1. Background

1.1. *Angiogenesis*

Angiogenesis describes the active biological process of blood vessel formation from pre-existing microvasculature (1, 2). In multi-celled organisms this is a highly organized and tightly regulated process that occurs normally during development, inflammation, and tissue repair. The importance of angiogenesis is reflected in the need of mammalian cells for oxygen and nutrients. Mammalian cells must be within a 200 μ M distance of blood vessels, which is the diffusion limit for oxygen (3). Thus the overall driving factor for angiogenesis is the requirement for oxygen and nutrients. The normal regulation of angiogenesis is mediated by the balance between pro- and anti-angiogenic factors that are released in the tissues and are influenced by local environmental factors.

1.2. *Angiogenesis and neoplasms*

In a neoplastic situation, the balances of these pro- and anti-angiogenic factors are generally skewed in favor of angiogenesis. In this setting, angiogenesis is generally a highly disorganized and loosely regulated process that is an absolute requirement for the continued growth of neoplasms (3). Further, there is a direct correlation between the extent of vascularization found in neoplasms and the potential for metastasis (4).

1.3. *Angiogenesis and chemokines*

There are a number of pro- and anti-angiogenic factors that have been described to date (3). The focus of this analysis will be on the chemokines interferon- γ -inducible protein (IP-10) and monokine induced by interferon- γ (MIG). Chemokines are a collection of cytokines that possess chemoattracting properties (for review see (5)). Chemokines are classified on the basis of the motif displayed by the first two cysteine residues present in the protein (CXC, CC, C, or CX3C), and they signal through G-protein coupled, seven-

transmembrane receptors. Initially identified for their influence on hemopoietic cell migration, chemokines are now known to influence a number of physiological and pathological process including angiogenesis and angiostasis (5).

IP-10 and MIG belong to a subset of the family of CXC chemokines (2) that bind the chemokine receptor CXCR3 (6). The CXC chemokine family can be further subdivided based on the presence or absence of a Glu-Leu-Arg or ELR motif at the NH₂ terminus of the chemokine. CXC chemokines that contain the ELR motif are potent promoters of angiogenesis whereas CXC chemokines that lack the ELR motif, as is the case for IP-10 and MIG, are potent inhibitors of angiogenesis (2).

2. Material and Methods

2.1. ODNs

ODN 1826 (TCCATGACGTTCTGACGTT; SEQ ID NO: 69)

2.2 Matrigel® - (BD)

Matrix solution is liquid at 4°C and solidifies at room temperature. When injected *in vivo* Matrigel solidifies to form a plug. Matrigel allows for the delivery of angiogenic promoters such as basic fibroblastic growth factor (bFGF) for the induction of angiogenesis. Plugs can then be removed to evaluate the level of angiogenesis as identified by the concentration of hemoglobin present. This system can be used to evaluate the anti-angiogenic potential of different compounds.

2.3 Hemoglobin quantification kit

Drabkin method reagent kit (Sigma)

2.4 Protein quantification

Protein quantification kit (BioRad)

2.5 Experimental design

For each group of 5 mice, the Matrigel was prepared as follows:

Group 1 - Matrigel alone.

3.5 mL of Matrigel

500 µL/mouse was injected subcutaneously (SC) right of center of the abdomen

Group 2 - Matrigel + bFGF (150ng/mL) + heparin (40 units/mL)

52.5 µL bFGF (10µg/mL)

23.2 µL heparin (6039 units/mL)

3.42 mL Matrigel

500 µL/mouse was injected SC right of center of the abdomen

Group 3 - Matrigel + bFGF (150 ng/mL) + heparin (40 units/mL) + oligo 1826 (1mg/mL)

52.5 µL bFGF (10µg/mL)

5 23.2 µL heparin (6039 units/mL)

233 µL oligo 1826 (15mg/mL)

3.19 mL Matrigel

500µL/mouse was injected SC right of center of the abdomen

Group 4 - Matrigel + bFGF (150ng/mL) + heparin (40 units/mL)

10 52.5 µL bFGF (10µg/mL)

23.2 µL heparin (6039 units/mL)

3.42 mL Matrigel

500 µL/mouse was injected SC right of center of the abdomen

This group received daily SC injections, for 6 days, of 100 µL of ODN 1826 (1mg/mL) on the opposite flank from the Matrigel plug.

15 **2.6 Determination of Hemoglobin and total protein content of Matrigel plugs**

On day 6 the animals were euthanised and the Matrigel plugs collected. The plugs were placed in 0.3 mL of sterile PBS and placed at 4°C over night to allow the Matrigel to liquify.

The hemoglobin and total protein content of the Matrigel plugs was determined using the methods described above. The hemoglobin content of the Matrigel plugs was expressed as (mg/mL)/mg of total protein.

20 **3. Preliminary Results**

When angiogenic factors were added to the Matrigel (Group 2), there was a significant increase in the amount of hemoglobin present in the Matrigel plug at 6 days when compared to Matrigel alone (Group 1) ($p < 0.05$). (See Figure 1.)

When CpG was included in the Matrigel plug along with the angiogenic factors (Group 3), there was a greater than 2 fold decrease in the amount of hemoglobin present in the Matrigel plug at 6 days when compared to the Matrigel containing the angiogenic factors (Group 2). (See Figure 1.)

When CpG was administered daily by subcutaneous injection, rather than present in the Matrigel plug, to the mouse in the flank opposite to the Matrigel plug which contained angiogenic factors (Group 4) there was no significant difference in the amount of hemoglobin

present in the Matrigel plug at 6 days when compared to Matrigel containing the angiogenic factors (Group 2). (See Figure 1.)

These preliminary results suggest that the inclusion of CpG ODN directly within the Matrigel (Group 3) had a negative influence on angiogenesis. Although daily delivery of CpG to the opposite flank from the Matrigel plug did not appear to influence angiogenesis, it is possible that CpG administered intravenously or subcutaneously in a region closer to the plug (and accordingly tumor mass) would manifest anti-angiogenic activity. CpG ODN may have to be present in the vicinity of active angiogenesis in order to have a negative influence.

4. References

1. Carmeliet, P. 2000. Mechanisms of angiogenesis and arteriogenesis. *Nat Med.* 6: 389-95.
2. Belperio, J. A., et al. 2000. CXC chemokines in angiogenesis. *J Leukoc Biol.* 68: 1-8.
3. Carmeliet, P., R. K. Jain. 2000. Angiogenesis in cancer and other diseases. *Nature.* 407: 249-57.
4. Zetter, B. R. 1998. Angiogenesis and tumor metastasis. *Annu Rev Med.* 49: 407-24.
5. Rossi, D., A. Zlotnik. 2000. The biology of chemokines and their receptors. *Annu Rev Immunol.* 18: 217-42.
6. Loetscher, M., et al. 1996. Chemokine receptor specific for IP10 and MIG: structure, function, and expression in activated T-lymphocytes. *J Exp Med.* 184: 963-9.
7. Coughlin, C. M., et al. 1998. Tumor cell responses to IFNgamma affect tumorigenicity and response to IL-12 therapy and antiangiogenesis. *Immunity.* 9: 25-34.
8. Strasly, M., et al. 2001. IL-12 inhibition of endothelial cell functions and angiogenesis depends on lymphocyte-endothelial cell cross-talk. *J Immunol.* 166: 3890-9.
9. Kanegane, C., et al. 1998. Contribution of the CXC chemokines IP-10 and Mig to the antitumor effects of IL-12. *J Leukoc Biol.* 64: 384-92.

Equivalents

It should be understood that the preceding is merely a detailed description of certain preferred embodiments. It therefore should be apparent to those of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. It is intended that the invention encompass all such modifications within the scope of the appended claims. All references, patents and patent applications and

publications that are cited or referred to in this application are incorporated in their entirety herein by reference.

I claim:

5

1001799-1301
TOTAL 66/007